



## Detoxification and Removal of Hexavalent Chromium in Aquatic Systems: Applications of Bioremediation

A.M.K.C.B. Aththanayake<sup>1</sup> , I.V.N. Rathnayake<sup>1</sup> ✉, M.P. Deeyamulla<sup>2</sup> 

<sup>1</sup>Department of Microbiology, Faculty of Science, University of Kelaniya, Kelaniya, GQ 11600, Sri Lanka.

<sup>2</sup>Department of Chemistry, Faculty of Science, University of Kelaniya, Kelaniya, GQ 11600, Sri Lanka.

Received: 01 Oct 2022; Revised: 01 Nov 2022; Accepted: 09 Nov 2022; Published online: 31 Dec 2022

### Abstract.

Chromium is a transition metal with a wide range of applications in leather tanning, textile, electroplating, stainless steel production, inorganic chemical production and wood preservation industries due to yellow colouration, corrosion resistance, higher melting-point and crystalline structure with ranging of oxidation states from 0 to +6. Trivalent and hexavalent chromium are the most abundant forms of chromium discharged into the aquatic environment by industries. It has been reported that hexavalent chromium is highly toxic than trivalent chromium due to the higher solubility, mobility and tendency to accumulate in higher trophic levels, which, therefore, become bioavailable and causes carcinogenic, mutagenic and teratogenic effects on most microorganisms and animals, growth inhibition, morphological and physiological changes and yield reductions in plants. Therefore, it is essential to detoxify the above hazardous pollutants up to permissible limits, which local and international authorities have legislated concerning its threat towards biotic components. Hexavalent chromium detoxification is possible to achieve using three methods i.e. physical, chemical and biological methods. These remediation processes can eliminate highly toxic hexavalent chromium or transform it into a less toxic form of trivalent chromium, completely or partially by adsorption and reduction. Biological remediation is considered a cost-effective and eco-friendly method compared to physical and chemical remediation. Further, many biological agents have been identified as agents that can tolerate the hexavalent chromium toxicity up to certain higher levels depending on the internal and external environmental factors, indicating different metal tolerance mechanisms that are assumed to be applied in metal remediation aspects. According to the testimonies of novel bioremediation studies, some hexavalent chromium tolerant organisms such as plants, bacteria, unicellular and multicellular fungi and algae are promising eco-friendly alternatives in detoxification and hexavalent chromium removal perspective. This article reviews the bioremediation approaches available for hexavalent chromium detoxification and removal and highlights the strengths and weaknesses of current bioremediation methods.

**Keywords:** Hexavalent chromium, toxicity, biological remediation, chemical remediation, physical remediation, biofilms

✉ Corresponding author, email: [vayanga@kln.ac.lk](mailto:vayanga@kln.ac.lk)

### Introduction

Chromium is a highly valued industrial raw material with a wide range of industrial applications such as pigment production for paints, inks and plastics, anti-corrosion coating production, stainless steel production, wood preservation and leather tannins which are discharged both trivalent chromium (Cr(III)) and hexavalent chromium (Cr(VI)) in higher quantities [1,2]. According to toxicological studies, Cr(VI) is 100 times more toxic than Cr(III) due to solubility, mobility and permeability in biota [3-7]. From the view of toxicology, prolonged exposure to Cr(VI) can lead to carcinogenic, mutagenic and teratogenic effects on animals which has been clinically proved, and morphological and physiological effects on plants, algae and other microorganisms [8-12]. The working community in chromium based industries, including chrome mining, has the highest potential for chromium poisoning [8,13-16]. Therefore, international authorities for the

occupational community, such as Occupational Safety and Health Administration (OSHA), has set the maximum limit for Cr(VI) exposures as similar to the conventional public health concerning local and international authorities; World Health Organization (WHO), United States Environmental Protection Agency (US EPA). The minimization and detoxification of Cr(VI) in industrial effluents can be achieved by Physical, Chemical, and biological methods. Among the above methods, biological remediation is considered the most cost-effective and environmentally friendly method. [17-20].

### Chemical nature and uses of Chromium.

Chromium is the 21<sup>st</sup> most abundant element in earth's crust, which belongs to d-block in the periodic table with a molar mass of 51.9961 g mol<sup>-1</sup> with a wide range of industrial applications based on chemical and physical properties such as inert nature, hardness, strength, high-temperature resistance and corrosion resistance etc. [21].



Cr(III) and Cr(VI) are the most stable and domain oxidation forms of chromium that exists in nature among the other oxidation forms of metallic chromium (Cr(0)), divalent chromium (Cr(II)), tetravalent chromium (Cr(IV)) and pentavalent chromium (Cr(V)).

The majority of chromium is used for metallurgical (67%) and refractories (18%), while the rest of 15 % are used for chromium induced chemical production, which is used for wood preservation, leather tanning, metal finishing, pigment production and textile industry as a raw material [2] (**Table 1**).

**Table 1** - Industrial applications of Cr(0), Cr(III) and Cr(VI).

Oxidation state	Industrial application	Reference
Cr (0)	Stainless steel production Alloy production Metal manufacturing	[2]
Cr (III)	Metal and alloy production Textile and leather tanning Copy machine toners Brick lining Chrome plating Catalysts production Paint production	[2,22]
Cr (VI)	Chrome plating Leather tanning Textile industry Copy machine toners Dye/paint pigment production Wood preservation High temperature battery production Metal finishing Catalyst production Stainless steel production Plastic production	[2,22,23]

### Toxicity of Chromium.

Among the most stable oxidation states of chromium in nature, Cr(III) is considered an essential micronutrient of higher organisms with less toxic effects due to lower solubility and impermeability [6,24]. Further, it has been reported that Cr(III) can assist in regulating the glucose level of the human body [25]. In contrast, Cr(VI) has been categorized as a carcinogenic agent by the USEPA and International Agency for Research on Cancer (IARC) due to high water solubility and mobility [26].

### Toxicity of Cr(VI) to humans.

Prolonged Cr(VI) exposes through breathing, ingesting, and skin contacts can cause nasal irritations, nasal perforations, skin irritations, skin ulcerations, skin

allergies, lung cancers, stomach upsets, convulsions, kidney and liver damages [2].

The working community in chromium-based industries (leather tanning, electroplating, mining and pigment production, etc.) has a high tendency to be affected by Cr(VI) toxicity. Cr(VI) can produce highly reactive hydroxyl radicals in blood vessels during the reduction into Cr(III), which can cause blood cell damages with organ degradations and cellular activity interruption by metal-DNA bindings [27]. Further it believes that Cr(VI) is responsible for causing teratogenic effects in human as it has proven with animal model trials [28].

### Toxicity of Cr(VI) to plants.

Chromium being a non-essential element it does not have a specific mechanism of uptake into plants. It is believed that, the plants use a passive process to uptake Cr(III) and an active process for uptake Cr(VI) with carriers competing with iron, sulphur and phosphorus [29]. Part of the Cr(VI) is taken up into plants after reducing into Cr(III) on the root surface, and the rest of the Cr(VI) is taken up by plants by dissolving in water and without reducing [29,30]. In the toxicological point of view, Cr(VI) affect plants both morphologically and physiologically. It has been found that, high concentrations of Cr(VI) affect the seed germination negatively due to the depressive effects on enzyme activity and sugar transport to embryo axes [31,32], It also reduces the root growth due to inhibition of water absorption [33], and shoot growth due to chromium transportation in aerial parts [31]. Toxicological Studies done elsewhere using *Oryza sativa*, *Acacia holosericea*, *Leucaena leucocephala* and *Albizia lebbek* and *Phaseolus vulgaris* reported that leaf area and biomass can be adversely affected by Cr(VI) [31,34].

Plant physiological studies revealed that Cr(VI) can lead to yield reduction by decreasing chlorophyll a, chlorophyll b and carotenoid pigments and affecting water and mineral transportation due to high oxidative potential [31,35].

### Toxicity of Cr(VI) to microorganisms.

Microorganisms are commonly exposed to many pollutants, including toxic metals, as they are widely dispersed in the environment, causing many toxic effects. Cr(VI) can become toxic to most bacterial strains causing cell enlargements, cell elongations and cell division inhibitions [23]. Cr(VI) can rapidly enter into the bacterial cytoplasm and reduces to lower oxidation states which are free radicals such as Cr(V), which leads to genotoxic effects by causing oxidative damages to DNA. Moreover, it has been found 400 – 800 µg of Cr(VI) can directly

interact with bacterial DNA causing frameshift mutations and base-pair replacements [36].

The Cr(VI) tolerance limits of bacteria have not clearly been defined as it can depend on several factors, including the type of the strain and physio-chemical conditions of the habitat, nature of the waste etc. Providing evidence to the above assumption a study of reported that 10 – 12 mg/L of Cr(VI) was adequate to inhibit most soil bacteria [22], while some strains in activated sludge can tolerate up to 80 mg/L of Cr(VI) [37]. Further, they have reported that Cr (VI) was able to stimulate bacterial growth up to 25 mg/L of Cr(VI).

Compared to bacteria, fungi are less sensitive to Cr(VI) due to the decreased uptake and production of antioxidants [38,39,39–41]. However, some studies describe that, Cr(VI) can cause genotoxic and mutagenic effects on several strains of fungi, including *Saccharomyces cerevisiae*, *Sclerotium rolfii* and *Pycnoporus sanguineus* leading to complex physiological changes and functional changes such as inhibition of oxygen uptake, induction of petite mutations and inducing mitochondrial functional damages [24,42–44]. Further studies on fungi have shown that effects of Chromium toxicity vary on the nature of carbon substrate [45].

Cr(VI) can affect PS II reaction centers of algae, which leads to inhibition of photosynthesis and cause significant morphological changes in some genera, including *Chlorella*, *Scenedesmas*, *Ulva*, *Isochrysis*, *Micrasterias*, and *Chlamydomonas* [36,46–51].

### Disposal and remediation process of Chromium wastes.

The chromium-based industries i.e., electroplating, tanning, water cooling, textile, wood preservation, alloy manufacturing, dye and pigment production discharge large quantities of contaminated chromium containing waste to soil, air and water annually. Considerable proportions of used chromium as a raw material and / or a reagent for industries including tannery (40%), chrome plating (35%), academic, research and industry laboratories (100%) discharge Cr(III) and Cr(VI) as effluents [52].

These chromium contaminated effluents should be remediated before discharging into the environment, due to toxicity of chromium to the environment and public health. Therefore, rules and regulations have been legislated and implemented by national and international authorized bodies such as WHO, US EPA, and national environmental acts of host countries for industrial wastewater and drinking water.

According to the US EPA and WHO standards maximum permissible level of Cr(VI) in drinking water and industrial wastewater have been legislated to 0.05 mg/L and 0.10 mg/L, respectively. Considering the health hazard to the occupational community in chromium-based industries, Occupational Safety and Health Administration (OSHA) has set the maximum limit for Cr(VI) compounds for 8-hour work shifts and 40-hour workweeks as 0.052 mg/L [23,53].

Based on these regulations, Cr(VI) contaminated wastes should be remediated before being discharged into the environment. Remediation of chromium containing waste can be carried out using three (03) methods i.e. chemical, physical and biological which are summarized in the **Table 2**.

### Chemical methods of Cr(VI) remediation.

Chemical reduction and photocatalysis are the most common chemical remediation methods that have been applied in chemical remediation processes. The chemical reduction uses reducing agents such as sulfur dioxide (SO<sub>2</sub>), calcium polysulfide (CaS<sub>5</sub>), ferrous sulfate (FeSO<sub>4</sub>), sodium metabisulfite (NaHSO<sub>3</sub>), sodium sulfite (Na<sub>2</sub>SO<sub>3</sub>), barium sulfite (BaSO<sub>3</sub>), hydrazine hydrate (N<sub>2</sub>H<sub>4</sub>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and, calcium carbonate (Na<sub>2</sub>CO<sub>3</sub>) [19,54–56]. Redox reactions of above mentioned reducing agents are kinetically slow at low Cr(VI) concentrations [57]. Therefore, it may require different methods to remediate residual Cr(VI), which are even higher than the discharge limits. Further, it has been found that this reduction process is also influenced by physical and chemical characteristics of the discharging sites (pH, conductivity, soil type and texture, presence of transition metals) [58,59].

Semiconductor based photocatalysis is a developing technology for toxic metal remediation such as Cr(VI), Hg(II), As(V), Cu(II), and Pb(II) [62]. This technology is more advantageous as there are no requirements for secondary disposal methods. Titania based photocatalysts such as TiO<sub>2</sub> and La<sub>2</sub>Ti<sub>2</sub>O<sub>7</sub> are extensively used for photocatalytic reduction of Cr(VI) in specific values [19,61]. But these Titania based photocatalysts cannot be applied practically to mass-scale commercial reactor systems due to high cost and operational disturbances due to sunlight irradiation and highly acidic conditions [62].

### Physical methods of Cr(VI) removal.

Physical remediation is achieved by techniques such as adsorption, electrolysis, ion exchange, membrane filtration and capping [19,63,64]. Adsorption is widely used for chromium removal in wastewater, consisting of

significant advantages such as low cost, profitability, availability, high efficiency, and minimum effort operation than other physio-chemical methods. A range of synthetic and natural adsorbents, including activated carbon, zeolite, chitosan, treated wastes, biological materials of coconut shell, wood husk, orange peel, hazelnut shell, sawdust, are used for Cr(VI) removal with a wide range of removal percentages under different pH values which are mostly laid on the extreme acidic range [65–69]. As some of the above adsorbents are freely available in nature, so that, adsorption is considered as one of the cost-effective methods of physical remediation. Membrane filtration technology is implemented with reverse osmosis, which is considered as one of the best available technology for removing all forms of chromium [70–72]. Literature shows that different membrane technology modifications to enhance Cr(VI) removal effectiveness, including micellar enhanced ultrafiltration, polymer inclusion membranes, ion exchange membranes and nanofiltration [73,74]. However, membrane technology is considered as a costly method with generating a large volume of concentrated liquid toxic wastes [72].

The acidic and basic ion exchange resins have also reported as effective Cr(VI) removal methods from chromium contaminated wastewater. A study of [75] have developed a complete Cr (VI) removal process from real wastewater by using a strongly basic synthetic Dowex 2-X4 resin without affecting pH. Further studies of [57] indicate 99.5% of Cr(VI) removal from “synthetic wastewater” using solvent impregnated resins which are acidic.

### **Biological methods of Cr(VI) removal.**

Remediation of chromium contaminated sources using biological agents including bacteria, fungi, algae, and plants play an important role in remediation approaches. Bacteria and fungi have shown efficient remediation agents than other agents. It has shown that organisms that can survive in a contaminated site may have the ability to remediate the contaminated site by themselves up to a certain level by transforming toxic pollutants into nontoxic forms [19,76–81]. This detoxification is achieved through biosorption, bioaccumulation and biotransformation.

Bioremediation is affected by several physio-chemical factors including, energy source (electron donors), electron acceptors, nutrients, pH, temperature and inhibitory substrates or metabolites [82]. Bioremediation of chromium is implemented in both in situ and ex situ

depending on the nature and requirements of the contaminated site [83,84].

Biological methods are considered as more advantageous from the economic and environmental point of view as they are cost-effective due to low installation and operational cost, eco-friendly with generating much less secondary pollutants, convenient and straightforward operation compared to physiochemical methods [63,85–87].

### **Bioremediation of Cr(VI) by bacteria.**

Bacterial bioremediation of Cr(VI) is explained in terms of chromium tolerance mechanisms such as biosorption and biotransformation/ bioreduction in both Gram-positive and Gram-negative strains [19]. During the bioreduction, highly toxic Cr(VI) is reduced into lesser toxic Cr (III) inside the bacterial cytoplasm, cell wall, or in both. The bacterial strains that can reduce Cr (VI) are usually named Chromium Reducing Bacteria (CRB). It is believed that Gram-positive CRB have a significant high tolerance to high Cr(VI) concentrations than gram-negative CRB [88].

According to previous studies, bacterial genera such as *Pseudomonas*, *Bacillus*, *Enterobacter*, *Deinococcus*, *Shewanella*, *Agrobacterium*, *Escherichia*, *Thermus*, *Microbacterium*, *Desulfovibrio*, *Deinococcus*, *Brucella*, and *Staphylococcus* have the potential to reduce Cr(VI) “directly” with enzymes and “indirectly” with metabolic end products [88–92]. It has also been reported that chromium tolerance and reduction are independent properties of bacteria, which means not all Cr(VI) resistant bacteria can reduce Cr(VI) into Cr(III) [88,93].

Bacterial Cr(VI) reduction is achieved under aerobic, anaerobic and both conditions [94]. Aerobic reduction is associated with soluble proteins and NADH as electron donors to enhance the reduction process, while anaerobic Cr(VI) reduction is associated with cell membrane bound reductase (flavin reductase, cytochromase, hydrogenases) and soluble reductase or both [95,96]. Bacterial bioreduction rate of Cr(VI) is influenced by initial cell density/ concentration, initial chromium concentration, initial pH, temperature, electron donors, oxyanions, salt concentration, presence of other heavy metals, metabolic inhibitors and oxidation-reduction potential of culture [96,97]. Further, the bacterial strains in the same species have different Cr(VI) tolerance and removal potentials depending on the level of the contaminants in the environment. This phenomena was evidenced in a comparative study carried out between uncontaminated and Cr(VI) polluted environments [98].

**Table 2.** Chemical, physical and biological method of Cr(VI) removal

Method	Technique	Mechanism	Type of contaminated source (Tested)	Cr, Cr (VI) removal percentage	Time	pH	Temp. (°C)	Reference
Chemical	Reduction	Cr(VI) reduction and adsorption by ED-RGO.	Synthetic wastewater	100%	24 hrs.	2.0	N/A	[120]
	Reduction	Cr(VI) reduction by Calcium polysulfide (CaS <sub>x</sub> ).	Contaminated ground water	90%	4 days	8 - 12.5	N/A	[121]
	Reduction and biosorption	Cr(VI) reduction and biosorption by chemically treated brown seaweed ( <i>Ecklonia</i> sp.).	Synthetic wastewater	100%	12 hrs.	2.0	25	[122]
	Reduction	Cr(VI) removal by chemically and electrochemically.	Synthetic wastewater	99.99%	10 min.	8.5 - 10.0	N/A	[123]
	Reduction	Cr(VI) reduction by Sodium corboxymethyl stabilized nanoscale zero valent iron.	Synthetic waste soil sample	80%	72 hrs.	4.73 - 7.36	N/A	[124]
	Reduction and coagulation	Cr(VI) removal by Ferrous sulfate (FeSO <sub>4</sub> ).	Spiked ground water	95%	46 hrs.	> 7.5	N/A	[125]
	Physical	Adsorption	Chromium Removal by fly ash.	Industrial waste	97.86%	12 hrs.	N/A	25
Adsorption		Cr(VI) removal by Ragi husk powder.	Synthetic wastewater	81.34%	2 hrs.	1.75	N/A	[127]
Adsorption		Cr(VI) removal by green algae and activated carbon.	Waste water	99.52%	2 hrs.	1.0	25	[110]
Adsorption		Cr(VI) removal by treated waste newspaper (TWNP).	Synthetic wastewater	64%	1 hrs.	3.0	25	[65]
Adsorption		Cr(VI) removal by green coconut shell.	Synthetic wastewater	95%	30 min.	6.5	28	[66]
Adsorption		Cr(VI) removal by agriculture wastes. Maize corncob. Cane bagasse. Jatropha oil cake.	Synthetic wastewater	62% 92% 97%	1 hrs.	2.0	30	[128]
Adsorption		Cr(VI) removal by <i>Mangifera indica</i> leaves.	Synthetic wastewater	91%	2 hrs.	2.0	30	[129]
Retention/ filtration		Cr(VI) removal by Aromatic polyimide thin film membrane.	Synthetic wastewater	77%	N/A	8.0	25	[73]
Adsorption		Cr(VI) removal by anion exchange resins.	Synthetic wastewater	99.4%	30 min.	3.0 - 5.0	25 60	[130]
Adsorption		Cr(VI) removal by hydrophobic resin.	Synthetic wastewater	99.5% 92%	24 hrs.	3.0	25	[131]
Adsorption		Cr(VI) removal by boiled rice husk.	Synthetic wastewater	71%	3 hrs.	2.0	27	[132]
Adsorption		Cr(VI) removal by formaldehyde treated rice husk.	Synthetic wastewater	76.5%	3 hrs.	2.0	27	[132]

Physical	Adsorption	Cr(VI) removal by modified montmorillonite clay nanocomposite.	Synthetic wastewater	99.9%	24 hrs.	2.0 - 6.6	25	[133]
	Adsorption	Cr(VI) removal by Fe- <sub>2</sub> O <sub>3</sub> / graphene adsorbents.	Synthetic wastewater	70.33%	N/A	3 - 4	25	[134]
	Adsorption	Cr(VI) removal by synthesized hydroxyapatite microfibrillated cellulose (CHA/MFC)	Synthetic wastewater	94%	5 min.	7 - 5	25	[135]
	Adsorption	Cr(VI) removal by Magnetite nanoparticles	Synthetic wastewater	66%	2 hrs.	3.0	25	[136]
	Adsorption	Cr(VI) removal by mixed waste tea and coffee ground	Synthetic wastewater	95%	3 hrs.	2.0	50 - 65	[137]
	Adsorption	Cr(VI) removal by natural adsorbents. Wool Olive cake Sawdust Pine needles Almond Coal Cactus	Synthetic wastewater	69.3% 47.1% 53.5% 42.9% 23.5% 23.6% 19.8%	2 hrs.	2.0	30	[138]
	Adsorption	Polypyrrole - montmorillonite clay composite	Synthetic wastewater	100%	24 hrs.	2.0	25	[139]
	Adsorption	Nanocomposite of ZnO with cotton stalks biochar	Synthetic wastewater	96.19%	1 hr.	2-4	25	[140]
	Filtration	Green emulsion liquid membrane	Synthetic wastewater	97-99%	0.5 hrs.	0.45	30	[71]
	Filtration	Green synthesized CuO nanoparticles	Synthetic wastewater	88.08%	2 hrs.	6.9	25	[70]
Biological	Reduction	Cr (VI) bioreduction by effluent bacteria <i>Staphylococcus cohnii</i>	Synthetic wastewater	90%	96 hrs.	7.2	37	[141]
	Reduction	Cr (VI) bioreduction by <i>Pseudomonas umsongensis</i>	Synthetic wastewater	93.9%	72 hrs.	7.0	30	[142]
	Reduction Adsorption	Cr (VI) bioreduction and biosorption by <i>Bacillus</i> sp.	Synthetic wastewater	97.04%	96 hrs.	7.0	37	[143]
	Reduction	Cr (VI) bioreduction by <i>Aeromonas hydrophila</i>	Synthetic wastewater	88%	72 hrs.	7.2	30	[144]
	Reduction	Cr(VI) reduction by <i>Bacillus thuringiensis</i>	Synthetic wastewater	86.42%	96 hrs.	7.0	35	[91]
	Reduction	Cr(VI) reduction by <i>Staphylococcus capitis</i>	Synthetic wastewater	97.34%	96 hrs.	7.0	35	[91]
	Reduction	Cr(VI) reduction by <i>Bacillus cereus</i>	Synthetic wastewater	98.5%	72 hrs.	7.1	26	[98]

Biological	Reduction Sorption	Cr(VI) reduction and sorption by <i>Enterobacter</i> sp.	Synthetic wastewater	99.1%	25 hrs.	6.0	45	[145]
	Reduction	Cr(VI) reduction by <i>Morganella morganii</i>	Synthetic wastewater	92%	48 hrs.	7.0	37	[146]
			Raw tannery effluent	90%	48 hrs.	7.0	37	
	Reduction Sorption	Cr(VI) reduction and sorption by <i>Stenotrophomonas rhizophila</i>	Synthetic wastewater	100%	28 hrs.	7.5	30	[147]
	Reduction	Cr(VI) reduction by <i>Cellulosimicrobium</i> sp.	Synthetic wastewater	100%	48 hrs.	7.0	30	[148]
	Reduction	Cr(VI) reduction by <i>Geobacter sulfurreducens</i>	Synthetic wastewater	99%	2hrs.	N/A	30	[149]
	Reduction	Cr(VI) reduction by <i>Pseudomonas aeruginosa</i>	Synthetic wastewater	93%	96 hrs.	7-8	30	[150]
	Sorption	Cr(VI) biosorption by <i>Shewanella putrefaciens</i>	Synthetic wastewater	85.68%	17 hrs.	8.0	38.44	[151]
	Reduction	Cr(VI) bioreduction by mixed bacterial consortium.	Synthetic wastewater	100%	120 hrs.	8.0	30	[152]
	Reduction Adsorption	Cr(VI) bioreduction and biosorption by <i>Corynebacterium paurometabolum</i> ,	Synthetic wastewater	55%	2 hrs.	3.0	30	[153]
	Reduction	Cr(VI) bioreduction by <i>Cellulosimicrobium funkei</i>	Synthetic wastewater	80.43%	120 hrs.	7.0	35	[154]
	Reduction	Cr(VI) bioreduction by <i>Pseudomonas stutzeri</i>	Synthetic wastewater	97%	24 hrs.	7.0	37	[155]
	Reduction	Cr(VI) bioreduction by <i>Acinetobacter baumannii</i>	Synthetic wastewater	99.58%	24 hrs.	8.0	37	[155]
	Reduction	Cr(VI) bioreduction by <i>Ochrobactrum</i> sp.	Synthetic wastewater	96.5%	N/A	7.0	30	[156]
	Adsorption	Cr(VI) biosorption by <i>Trichoderma</i> sp.	Synthetic wastewater	97.39%	2 hrs.	5.5	25	[103]
	Reduction Adsorption	Cr(VI) biosorption and bioreduction by <i>Paecilomyces lilacinus</i>	Synthetic wastewater	100%	120 hrs.	5.5	25	[157]
	Adsorption	Cr(VI) biosorption by <i>Phanerochaete chrysosporium</i>	Synthetic wastewater	99.7%	72 hrs.	7.0	40	[158]
	Adsorption	Cr(VI) biosorption by <i>Pleurotus ostreatus</i>	Synthetic wastewater	80%	12 hrs.	2.0 – 11.0	65	[159]
	Adsorption	Cr(VI) adsorption by Cationic surfactant-modified, <i>Kazachstania yasuniensis</i> <i>Kodamaea transpacifica</i> <i>Saturnispora quitensis</i> <i>Saccharomyces cerevisiae</i>	Synthetic wastewater	80.70% 85.80% 85.40% 75.80%	4 hrs.	4.5	25	[105]

	Adsorption	Cr(VI) adsorption by a hydroxyl-functionalized magnetic <i>Aspergillus niger</i> nanocomposite	Synthetic wastewater	64.91%	4 hrs.	5.0	50	[160]
Biological	Adsorption	Cr (VI) adsorption by <i>Chlorella sorokiniana</i> .	Synthetic waste water	99.68%	72 hrs.	7.0	40	[161]
	Reduction Adsorption	Cr (VI) removal by green algal strain <i>Cladophora albida</i>	Synthetic waste water industrial waste water	100%	120 hrs.	0.5	25	[162]
	Adsorption	Cr(VI) removal by marine algae <i>Turbinaria ornata</i>	Synthetic wastewater	95.25%	3.5 hrs.	4.7	33.6	[163]

N/A – Not applicable

**Table 3.** Microorganisms and substrates used in biofilm formation for bioremediation of Cr(VI)

Organism	Adhesion substrate	Cr (VI) removal percentage (%)	pH	Temp. (°C)	Time	Reference
<i>Arthrobacter viscosus</i>	Granular activated carbon	99.9%	5 – 5.5	28	30 days	[190]
<i>Pseudomonas</i> sp. <i>Bacillus</i> sp. <i>Azotobacter</i> sp. <i>Acremonium</i> sp.	Glass wool	90%	5.6-6.1	30	10 days	[172]
<i>Streptomyces</i> strain CG252	Glass bead	100%	N/A	30	48 – 72 hrs.	[191]
<i>Arthrobacter</i> sp.	Gravel packed bed reactors	100%	N/A	30	26 hrs.	[192]
<i>Morganella morgani</i> STB5	Polystyrene Polysulfone	99.47% 90.78%	7.0	30	72 hrs.	[193]
<i>Arthrobacter</i> sp. SUK 1205	Glass beads	100%	7.0	37	96 hrs.	[194]
<i>Halomonas</i> sp.	Pumic particle stones	94.5%	6.5	28	48 hrs.	[195]
<i>Bacillus subtilis</i> <i>Escherichia coil</i> <i>Acinetobacter junii</i>	Alginate bead	97.84%	7.0	25	7 hrs.	[196]
<i>Wickerhamomyces anomalus</i>	Wood husk	92.5%	3.72	30	N/A	[171]
<i>Acinetobacter haemolyticus</i>	Wood husk	97%	7.0	25	72 hrs.	[69]
<i>Streptococcus salivarius</i> <i>Pseudomonas fluorescens</i> LB 300	Stainless steel AISI 316L Glass beads	42% 100%	N/A 6.8-7.0	37 30	72 hrs. 8 days	[197] [198]
<i>Cellulosimicrobium</i> sp.	PVC Rubber tubing Sand Small stone	99.5% 90.0% 96% 88.4%	N/A	25	11 days	[199]
<i>Escherichia coli</i>	Kaolin	100%	4.6-5.1	37	10 days	[200]
<i>Nostoc</i> sp.	Polystyrene	86.49%	7.0	25	7 days	[201]
<i>Shewanella xiamenensis</i>	Zeolite	100%	3.0	22-25	35 days	[202]
<i>Cunninghamella elegans</i>	Stainless steel compression springs	98.6%	7.0-3.0	28	40 hrs.	[203]
<i>Arthrobacter</i> sp. SUK 1201	Glass beads	100%	7.0	37	3 days	[204]
<i>Lysinibacillus sphaericus</i> RTA-01	Glass slide	82.8%	5.2	37	72 hrs.	[205]
<i>Ochrobactrum pseudintermedium</i> ADV31	Polyurethane foam	82%	7.0	45	5 days	[206]

N/A – Not applicable

### Bioremediation of Cr(VI) by fungi.

Similar to the bacterial remediation of toxic metals, some fungal strains have been investigated for bioremediation with the same metal removal techniques used with bacteria, i.e. biosorption, bioaccumulation and biotransformation/ bioreduction.

Fungi *Aspergillus* sp. was reported to remove chromium through bioreduction from contaminated effluents [99]. This study also indicates 65% of chromium removal from tannery effluent and 85% of Cr(VI) removal from the synthetic medium at pH 6 within 07 days. The similar bioreduction of Cr(VI) has been also reported by [19,100–102] with *Hypocrea tawa*, *Trichoderma inhamatum* and isolated Yeast strains. However, the Cr(VI) reduction capability of fungal strains can be changed with the initial Cr(VI) concentration and initial biomass of the strain.

*Fusarium oxysporum* and *Trichoderma* sp. have shown to adsorb Cr(VI) on to their cell surface by forming chemical bonds with cell surface proteins with analytically verified evidence using FT-IR spectrum [19,103]. Comparison of Cr(VI) removal in synthetic and raw wastes was found to be 77% and 85% of Cr(VI) removal respectively, with 200-1000 mg/L of initial Cr(VI) concentrations using immobilized Baker's yeast strain (*Saccharomyces cerevisiae*) in Biomass/Polymer Matrices Beads (BPMM) through biosorption [104]. A study comparing Cr(VI) biosorption by native Ecuadorian yeast species, reported that *Kazachstania yasuniensis*, *Kodamaea transpacificica*, and *Saturnispora quitensis* have the ability to remove Cr(VI). Furthermore, they have reported that efficient Cr(VI) removal can be achieved by inducing belzalkonium chloride (BZK) to cell surface as a chemical modification to the applying bio agent [105].

### Bioremediation of Cr(VI) by algae.

It is evident that both freshwater and marine algal species such as *Cladophora* sp., *Selenastrum* sp., *Spirogyra* sp., *Ceramium* sp, *Chlorella* sp. and *Ulva* sp. can be used to remediate chromium contaminated wastewater by applying as cultures or incorporating with other physio-chemical methods following biosorption and bioreduction [19,106–109]. Unlike other organisms, algae have been used in both living and non-living forms for Cr(VI) remediation. Introducing an efficient Cr(VI) removal method [110] reports that dried *Ulva lactuca* incorporated into activated carbon can be used to remediate highly acidic and halophilic wastewater. *Chlorella* sp. has been used in most Cr(VI) algal remediation bioreactors as it is widely dispersed in the aquatic environment with higher Cr(VI) tolerance [109,111–114]. Constructing a hybrid remediation system

[115] has introduced efficient and reusable alumina hollow fibers immobilized with TiO<sub>2</sub> and *Chlorella vulgaris* cells. Additionally, this hybrid system has been achieved greater than 90% of Cr(VI) removal after five sequential reuses.

However, similar to bacterial bioremediation, algal Cr(VI) bioremediation is influenced by physio-chemical parameters including pH, temperature, initial biomass, initial Cr(VI) concentration, light intensity, contact time of cells and wastes of the treatment process bioremediation [19,116].

### Bioremediation of Cr(VI) by plants.

Limited studies have reported that the Cr(VI) detoxification and removal potential of plants compared to the other biological agents. Green plants detoxify many pollutants using various mechanisms followed by uptake, known as phytoremediation [117]. Plants can either store heavy metals in roots or partially translocate to shoot through the xylem after getting diffused into the root system. Further, it has been reported that upward translocation of heavy metals is retarded by the cation exchange process in plant tissues and leads to considerable heavy metal accumulation in roots compared to axial parts of plants. In the point of view of Cr(VI) and other chromium forms, this phenomenon has been reported else ware using *Phragmites australis*, *Ailantus altissima* and *Salix viminalis* [118]. This may be due to the encapsulation in vacuoles of root cells based on natural counteraction of plants against chromium toxicity [119]. According to observations over 360 days, a study suggests that *Salix viminalis* used for large scale phytoremediation application for removal of Cr(VI) and other chromium forms from contaminated sources as *S.viminalis* were removed 70% of total chromium and 90% of Cr(VI) removal with indicating higher translocation capacity [118].

In vitro study of *Nopalea cochenillifera* found that it has the potential to accumulate a wide range of Cr(VI) (600 – 26,000 mg/ Kg) from the growth medium. As *N. cochenillifera* is a non-consuming plant for diets, the risk of bioaccumulation can be avoided in the ecosystem. Furthermore, the above study has also reported plant species with different chromium accumulation potentials including *Gynura pseudochina*, *Brassica napus*, *Prosopis juliflora*, *Leersia hexandra*, *Urtica dioica*, *Salix matsudana*, *Brassica napus*, *Helianthus annuus*, *Lycopersicon lycopersicum* and *Saponaria officinalis* perhaps considered for the phytoremediation [117].

## Biofilms

An aggregated community of prokaryotic and eukaryotic microorganisms adhering to substance/matrix surface and submerged/embedded in self-produced extracellular polymeric substances (EPS) is termed a "biofilm" [164,165]. These aggregates are omnipresent in the biosphere, including soil, water, plant and animal tissues, abiotic substances like pipelines, ship hulls and filters. These biofilms can be developed in solid-water, water-air and solid-air interfaces with composing EPS, multivalent cations, biogenic particles, colloidal and dissolved compounds [166]. Biofilms are comprised of both single and multiple microbial species. Among them, multiple species biofilms are the most dispersed biofilm type in the environment [167].

Biofilm formation is a subsequent process that consists of 03 main steps; surface attachment, biofilm maturation and dispersal [165]. In surface attachment, microbial cells undergo reversible attachment at cell poles by involving cell appendages (flagella, pilli and fimbriae) followed by irreversible attachment. After the reversible attachment stage, microbial cells can be adapted to biofilm lifestyle or left the matrix. During the irreversible attachment, stage cells adhere to the matrix by EPS and surface proteins (Sad B and Lap A).

Biofilm maturation starts after the irreversible attachment with developing microcolonies. At this stage, previous microbial cells are assembled and proliferated along with producing EPS. Further studies explain that biofilm structure, including thickness and cell density, is dynamically changed according to environmental conditions such as temperature, presence of oxygen, pH and amounts of nutrients [164,165].

The immobilized microbial cells are transferred back to planktonic growth as the final stage of the biofilm lifecycle and as an initial step of a new forming biofilm. This dispersal can be happened "actively" by cell motility and EPS degradation or "passively" by external physical forces.

## Environmental applications of biofilms

Even though free-living microorganisms are capable of bioremediating polluted environments, the remediation process can be disrupted due to high concentrated toxic compounds, availability of nutrients and environmental stress. Applying sessile or floating biofilms for remediation is highly advantageous as biofilm communities have higher tolerance towards environmental stress, including lack of nutrient availability, high concentrated chemical exposures, pH

and temperature fluctuations, lack of moisture content than free-living microorganisms [168].

When selecting biofilms for remediation purposes, several factors must be considered: the capability to tolerate environmental stress, exchange of genetic materials, growth rates, metabolic diversity, and symbiotic relationships. Based on above factors bacterial, algal and fungal biofilms are used in bioreactors to remediate contaminated sources by a wide range of pollutants including, organic pollutants (polyaromatic hydrocarbons, chlorinated aromatic compounds, aromatic amine compounds, polyethylene and polythene), heavy metals (Cu, Zn, Cd, Ni, As, Fe, Hg, Mn), inorganic pollutants (nitrate ions and synthetic dyes) contaminates which can adversely affect the eco-systems [168-170].

## Cr(VI) remediation by biofilms

Cr(VI) bioremediation is achieved using bacterial, algal and fungal single species and multi-species biofilms growing on either natural or artificial substrates through bioremediation and biosorption techniques [171-174].

Biofilms are more effective for Cr(VI) remediation than planktonic cells. The study investigating *Streptomyces* sp. strain CG252 and *Pseudomonas aeruginosa* A2Chr, respectively, reported evidence for the above phenomena [102,175]. Another study using three (03) different biosorbents including lyophilized *Escherichia coli* AUS 7 cells, granulated activated carbon (GAC) and biofilm of *Escherichia coli* AUS 7 on GAC exhibited that biofilms are able to achieve a higher adsorption of Cr(VI) than GAC and lyophilized cells in according to Langmuir and Freundlich isotherm models from aqueous solutions under acidic conditions [176]. However, contradictory results were also reported elsewhere, that the planktonic cells have more significant potential for Cr(VI) reduction than biofilms based on their study of *Bacillus subtilis* ATCC-6633. Furthermore above study revealed that, biofilm debris are susceptible to immobilized reduced Cr(III) ions completely [177].

Compared to bacteria, reports on Cr(VI) remediation by fungal biofilms are scarce in the literature. Immobilized cells of *Aspergillus niger*, *Coriolus versicolor*, *Saccharomyces cerevisiae*, and *Lentinus sajorcaju* have been used in sorption and reduction techniques [99,178-180]. Moreover, immobilized algal cells on different matrixes have been used for Cr(VI) remediation by sorption of metal ions to the cell wall components [181-185].

Algal-bacterial biofilms/consortia have also been used for Cr(VI) bioremediation. Further, it is reported that these consortia have symbiotic effects on each other by

supplying each other's nutritional needs such as O<sub>2</sub> for aerobic bacteria by algae and CO<sub>2</sub> for algae by bacteria during the remediation process. Therefore, algal-bacterial consortia are termed as a self-sustaining system [186–188]. The algal-bacterial system has the potential to remove higher Cr(VI) contents such as 100 mg/L, 75 mg/L and 50 mg/L providing a carbon source to the mixed consortium of chromium reducing bacterial (CRB) cultures of *Escherichia coli*, *Bacillus thermoamylovorans* and *Citrobacter sedakii* from the algal strain of *Chlamdomonas reinhartii*. Furthermore, the above study suggests that the algal-biofilm consortia as a cost-effective method that prevents cost of carbon sources as it fulfils by algae even though algal-bacterial consortia takes a longer time duration for the Cr(VI) removal [189]. **Table 3** illustrates some of the microorganisms and substrates used in bioremediation of Cr(VI).

### Limitations and remedial actions of the current bioremediation methods in Cr(VI) removal

The notable limitations of biological methods available for Cr(VI) removal have been identified including, varying Cr(VI) tolerance and removal levels environmental conditions and nutritional requirements of biological components used, toxic substances present in wastewater which can interfere with the biological components, disposal of the accumulated Cr in the biological component, and practical difficulties in extrapolating bench/pilot-scale to full-scale field application [207–209]. Customized solutions need to be sought by assessing the remediation requirements at individual level, because the environmental conditions and the nutritional requirements of the biological component vary depending on the contaminated site. Moreover, to overcome the Cr disposal after bioremediation, it is possible to percolate the biotransformed Cr(III) through reduction at low pH conditions and tend to be reoxidised to Cr(VI) in the presence of manganese oxide and chlorine in treated effluent or discharging environment [210,211]. In order to prevent the discharge of higher amounts of chromium and to enhance the sorption capacity of biomasses, the metal desorption process should be followed. This desorption can be done by acid digestions [212–215] and alkaline treatments [216–220] as a hybrid Cr(VI) remediation process, which has many benefits such as reduction of generation of secondary pollutants and recovery of valuable metals. These recovered Cr(VI) and Cr(III) can be applied for tannery and chromium-based chemical production as raw materials [221].

## Conclusion

The wide industrial and research application of Chromium followed by emitting considerable amounts of Cr(VI), coupled with the fact that it leads to serious problems to all components of the ecosystem. Therefore, it has been legislated to remediate Cr(VI) contaminated effluents by national and international authorities before it being discharged to the environment. This remediation is carried out by chemical, physical and biological methods. Biological remediation is considered as the most environment-friendly and cost-effective method rather than chemical and physical remediation. However, considering the limitations of the current bioremediation processes, hybrid remediation processes combining the bioremediation with other chemical and physical methods are being used for the effective remediation of Cr(VI) in aquatic systems.

## Authors Contribution

The authors confirm contribution to the paper as follows: study conception and design: A.M.K.C.B. Aththanayake, I.V.N. Rathnayake, and M.P. Deeyamulla; draft manuscript preparation: A.M.K.C.M. Aththanayake; Review, and editing the final draft: A.M.K.C.B. Aththanayake, I.V.N. Rathnayake; All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no any competing interest.

## Acknowledgement

This research was supported by the National Research Council, Sri Lanka Investor Driven Research Grant No. 18-083.

## References

1. Park D, Yun Y-S, Park JM. Use of dead fungal biomass for the detoxification of hexavalent chromium: screening and kinetics. *Process Biochem* [Internet]. 2005 Jun [cited 2020 Apr 9];40(7):2559–65. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0032959205000038>
2. Saha R, Nandi R, Saha B. Sources and toxicity of hexavalent chromium. *J Coord Chem* [Internet]. 2011 May 20 [cited 2020 Apr 9];64(10):1782–806. Available from: <https://www.tandfonline.com/doi/full/10.1080/00958972.2011.583646>
3. Ashraf A, Bibi I, Niazi NK, Ok YS, Murtaza G, Shahid M, et al. Chromium(VI) sorption efficiency of acid-activated banana peel over organo-montmorillonite in aqueous solutions. *Int J Phytoremediation* [Internet]. 2017 Jul 3 [cited 2021 Sep 18];19(7):605–13. Available from: <https://www.tandfonline.com/doi/full/10.1080/15226514.2016.1256372>
4. Choppala G, Bolan N, Lamb D, Kunhikrishnan A. Comparative Sorption and Mobility of Cr(III) and Cr(VI) Species in a Range of Soils: Implications to Bioavailability. *Water Air Soil Pollut* [Internet]. 2013 Dec [cited 2021 Jul 11];224(12):1699. Available from: <http://link.springer.com/10.1007/s11270-013-1699-6>
5. Laccalle RG, Garbisu C, Becerril JM. Effects of the application of an organic amendment and nanoscale zero-valent iron particles on



- soil Cr(VI) remediation. *Environ Sci Pollut Res* [Internet]. 2020 Sep [cited 2021 Sep 18];27(25):31726–36. Available from: <https://link.springer.com/10.1007/s11356-020-09449-x>
6. Megharaj M, Avudainayagam S, Naidu R. Toxicity of Hexavalent Chromium and Its Reduction by Bacteria Isolated from Soil Contaminated with Tannery Waste. *Curr Microbiol* [Internet]. 2003 Jul 1 [cited 2019 Feb 27];47(1):51–4. Available from: <http://link.springer.com/10.1007/s00284-002-3889-0>
  7. Xia S, Song Z, Jeyakumar P, Shaheen SM, Rinklebe J, Ok YS, et al. A critical review on bioremediation technologies for Cr(VI)-contaminated soils and wastewater. *Crit Rev Environ Sci Technol* [Internet]. 2019 Jun 18 [cited 2020 May 5];49(12):1027–78. Available from: <https://www.tandfonline.com/doi/full/10.1080/10643389.2018.1564526>
  8. Anttila S, Boffetta P, editors. *Occupational Cancers* [Internet]. Cham: Springer International Publishing; 2020 [cited 2021 Sep 20]. Available from: <http://link.springer.com/10.1007/978-3-030-30766-0>
  9. Chen QY, Murphy A, Sun H, Costa M. Molecular and epigenetic mechanisms of Cr(VI)-induced carcinogenesis. *Toxicol Appl Pharmacol* [Internet]. 2019 Aug [cited 2021 Jul 12];377:114636. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0041008X19302443>
  10. Chervona Y, Costa M. Hexavalent Chromium and Cancer. In: Kretsinger RH, Uversky VN, Permyakov EA, editors. *Encyclopedia of Metalloproteins* [Internet]. New York, NY: Springer New York; 2013 [cited 2021 Sep 20]. p. 969–75. Available from: [http://link.springer.com/10.1007/978-1-4614-1533-6\\_10](http://link.springer.com/10.1007/978-1-4614-1533-6_10)
  11. Prado FE, Hilal M, Chocobar-Ponce S, Pagano E, Rosa M, Prado C. Chromium and the Plant. In: *Plant Metal Interaction* [Internet]. Elsevier; 2016 [cited 2021 Sep 20]. p. 149–77. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780128031582000060>
  12. Wani PA, Wani JA, Wahid S. Recent advances in the mechanism of detoxification of genotoxic and cytotoxic Cr (VI) by microbes. *J Environ Chem Eng* [Internet]. 2018 Aug [cited 2021 Sep 20];6(4):3798–807. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2213343718302884>
  13. Oginawati K, Susetyo SH, Rosalyn FA, Kurniawan SB, Abdullah SRS. Risk analysis of inhaled hexavalent chromium (Cr6+) exposure on blacksmiths from industrial area. *Environ Sci Pollut Res* [Internet]. 2021 Mar [cited 2021 Sep 20];28(11):14000–8. Available from: <http://link.springer.com/10.1007/s11356-020-11590-6>
  14. Wild P, Bourgkard E, Paris C. Lung Cancer and Exposure to Metals: The Epidemiological Evidence. In: Verma M, editor. *Cancer Epidemiology* [Internet]. Totowa, NJ: Humana Press; 2009 [cited 2021 Sep 20]. p. 139–67. (Walker JM, editor. *Methods in Molecular Biology*; vol. 472). Available from: [http://link.springer.com/10.1007/978-1-60327-492-0\\_6](http://link.springer.com/10.1007/978-1-60327-492-0_6)
  15. Yoshinaga M, Ninomiya H, Al Hossain MMA, Sudo M, Akhand AA, Ahsan N, et al. A comprehensive study including monitoring, assessment of health effects and development of a remediation method for chromium pollution. *Chemosphere* [Internet]. 2018 Jun [cited 2021 Sep 20];201:667–75. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653518304314>
  16. Zhang X-H, Zhang X, Wang X-C, Jin L-F, Yang Z-P, Jiang C-X, et al. Chronic occupational exposure to hexavalent chromium causes DNA damage in electroplating workers. *BMC Public Health* [Internet]. 2011 Dec [cited 2021 Sep 20];11(1):224. Available from: <https://bmcpubhealth.biomedcentral.com/articles/10.1186/1471-2458-11-224>
  17. Haq I, Kalamdhad AS, editors. *Emerging Treatment Technologies for Waste Management* [Internet]. Singapore: Springer Singapore; 2021 [cited 2021 Sep 20]. Available from: <https://link.springer.com/10.1007/978-981-16-2015-7>
  18. He C, Gu L, Xu Z, He H, Fu G, Han F, et al. Cleaning chromium pollution in aquatic environments by bioremediation, photocatalytic remediation, electrochemical remediation and coupled remediation systems. *Environ Chem Lett* [Internet]. 2020 May [cited 2021 Sep 20];18(3):561–76. Available from: <http://link.springer.com/10.1007/s10311-019-00960-3>
  19. Jobby R, Jha P, Yadav AK, Desai N. Biosorption and biotransformation of hexavalent chromium [Cr(VI)]: A comprehensive review. *Chemosphere* [Internet]. 2018 Sep [cited 2020 Apr 9];207:255–66. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653518308981>
  20. Karimi-Maleh H, Ayati A, Ghanbari S, Orooji Y, Tanhaei B, Karimi F, et al. Recent advances in removal techniques of Cr(VI) toxic ion from aqueous solution: A comprehensive review. *J Mol Liq* [Internet]. 2021 May [cited 2021 Sep 20];329:115062. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0167732220373049>
  21. Barnhart J. Occurrences, Uses, and Properties of Chromium. *Regul Toxicol Pharmacol* [Internet]. 1997 Aug [cited 2020 Apr 9];26(1):S3–7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0273230097911326>
  22. Losi ME, Amrhein C, Frankenberger WT. Environmental Biochemistry of Chromium. In: Ware GW, editor. *Reviews of Environmental Contamination and Toxicology* [Internet]. New York, NY: Springer New York; 1994 [cited 2020 Apr 10]. p. 91–121. (Reviews of Environmental Contamination and Toxicology; vol. 136). Available from: [http://link.springer.com/10.1007/978-1-4612-2656-7\\_3](http://link.springer.com/10.1007/978-1-4612-2656-7_3)
  23. Mishra S, Bharagava RN. Toxic and genotoxic effects of hexavalent chromium in environment and its bioremediation strategies. *J Environ Sci Health Part C* [Internet]. 2016 Jan 2 [cited 2021 Jul 10];34(1):1–32. Available from: <http://www.tandfonline.com/doi/full/10.1080/10590501.2015.1096883>
  24. Cervantes C, Campos-García J, Devars S, Gutiérrez-Corona F, Loza-Tavera H, Torres-Guzmán JC, et al. Interactions of chromium with microorganisms and plants. *FEMS Microbiol Rev* [Internet]. 2001 May [cited 2020 Apr 9];25(3):335–47. Available from: <https://academic.oup.com/femsre/article-lookup/doi/10.1111/j.1574-6976.2001.tb00581.x>
  25. Costa M, Klein CB. Toxicity and Carcinogenicity of Chromium Compounds in Humans. *Crit Rev Toxicol* [Internet]. 2006 Jan [cited 2020 Apr 9];36(2):155–63. Available from: <http://www.tandfonline.com/doi/full/10.1080/10408440500534032>
  26. McLean JE, McNeill LS, Edwards MA, Parks JL. Hexavalent chromium review, part 1: Health effects, regulations, and analysis. *J - Am Water Works Assoc* [Internet]. 2012 Jun [cited 2020 Apr 9];104(6):E348–57. Available from: <http://doi.wiley.com/10.5942/jawwa.2012.104.0091>
  27. Kim HS, Kim YJ, Seo YR. An Overview of Carcinogenic Heavy Metal: Molecular Toxicity Mechanism and Prevention. *J Cancer Prev* [Internet]. 2015 Dec 30 [cited 2020 Apr 9];20(4):232–40. Available from: <http://www.jcpjournal.org/journal/view.html?doi=10.15430/JCP.2015.20.4.232>
  28. Baruthio F. Toxic effects of chromium and its compounds. *Biol Trace Elem Res* [Internet]. 1992 Jan [cited 2020 Apr 9];32(1–3):145–53. Available from: <http://link.springer.com/10.1007/BF02784599>
  29. Oliveira H. Chromium as an Environmental Pollutant: Insights on Induced Plant Toxicity. *J Bot* [Internet]. 2012 May 20 [cited 2020 Apr 9];2012:1–8. Available from: <https://www.hindawi.com/archive/2012/375843/>
  30. Zayed A, Lytle CM, Qian J-H, Terry N. Chromium accumulation, translocation and chemical speciation in vegetable crops. *Planta* [Internet]. 1998 Aug 6 [cited 2020 Apr 9];206(2):293–9. Available from: <http://link.springer.com/10.1007/s004250050403>
  31. Shanker A, Cervantes C, Lozatarvera H, Avudainayagam S. Chromium toxicity in plants. *Environ Int* [Internet]. 2005 Jul [cited 2020 Apr 9];31(5):739–53. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0160412005000231>
  32. Zeid IM. Responses of *Phaseolus Vulgaris* Chromium and Cobalt Treatments. *Biol Plant* [Internet]. 2001 [cited 2020 Apr 9];44(1):111–5. Available from: <http://link.springer.com/10.1023/A:1017934708402>
  33. Barceló J, Poschenrieder C, Gunsé B. Water Relations of Chromium VI Treated Bush Bean Plants (*Phaseolus vulgaris* L. cv. Contender)



- under both Normal and Water Stress Conditions. J Exp Bot [Internet]. 1986 [cited 2020 Apr 9];37(2):178-87. Available from: <https://academic.oup.com/jxb/article-lookup/doi/10.1093/jxb/37.2.178>
34. Karunyal S, Renuga G, Kailash P. Effects of tannery effluent on seed germination, leaf area, biomass and mineral content of some plants. Bioresour Technol [Internet]. 1994 Jan [cited 2020 Apr 9];47(3):215-8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/096085249490183X>
  35. Nichols PB, Couch JD, Al-Hamdani SH. Selected physiological responses of *Salvinia minima* to different chromium concentrations. Aquat Bot [Internet]. 2000 Dec [cited 2020 Apr 9];68(4):313-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304377000001285>
  36. Laxmi V, Kaushik G. Toxicity of Hexavalent Chromium in Environment, Health Threats, and Its Bioremediation and Detoxification from Tannery Wastewater for Environmental Safety. In: Saxena G, Bharagava RN, editors. Bioremediation of Industrial Waste for Environmental Safety [Internet]. Singapore: Springer Singapore; 2020 [cited 2020 Apr 11]. p. 223-43. Available from: [http://link.springer.com/10.1007/978-981-13-1891-7\\_11](http://link.springer.com/10.1007/978-981-13-1891-7_11)
  37. Sepideh S, S B, C S, B K. A Study of Toxic Dosage of Combined Selenium and Hexavalent Chromium on Activated Sludge Bacteria. Int J Water Wastewater Treat [Internet]. 2019 [cited 2020 Apr 11];5(1). Available from: <https://www.sciforschenonline.org/journals/water-and-waste/IJWWT161.php>
  38. Ahemad M. Bacterial mechanisms for Cr(VI) resistance and reduction: an overview and recent advances. Folia Microbiol (Praha) [Internet]. 2014 Jul [cited 2021 Jul 12];59(4):321-32. Available from: <http://link.springer.com/10.1007/s12223-014-0304-8>
  39. Kumar V, Dwivedi SK. Hexavalent chromium stress response, reduction capability and bioremediation potential of *Trichoderma* sp. isolated from electroplating wastewater. Ecotoxicol Environ Saf [Internet]. 2019 Dec [cited 2021 Sep 7];185:109734. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0147651319310656>
  40. Poljsak B, Pócsi I, Raspor P, Pesti M. Interference of chromium with biological systems in yeasts and fungi: a review: Effects of chromium on yeast and fungi. J Basic Microbiol [Internet]. 2010 Feb [cited 2020 Apr 10];50(1):21-36. Available from: <http://doi.wiley.com/10.1002/jobm.200900170>
  41. Vajpai S, Taylor PE, Adholeya A, Leigh Ackland M. Chromium tolerance and accumulation in *Aspergillus flavus* isolated from tannery effluent. J Basic Microbiol [Internet]. 2020 Jan [cited 2021 Sep 7];60(1):58-71. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/jobm.201900389>
  42. Feng M, Yin H, Peng H, Liu Z, Lu G, Dang Z. Hexavalent chromium induced oxidative stress and apoptosis in *Pycnoporus sanguineus*. Environ Pollut Barking Essex 1987. 2017 May 18;228:128-39.
  43. Kharab P, Singh I. Genotoxic effects of potassium dichromate, sodium arsenite, cobalt chloride and lead nitrate in diploid yeast. Mutat Res Toxicol [Internet]. 1985 Mar [cited 2020 Apr 10];155(3):117-20. Available from: <https://linkinghub.elsevier.com/retrieve/pii/0165121885901284>
  44. Shahid M, Shamshad S, Rafiq M, Khalid S, Bibi I, Niazi NK, et al. Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: A review. Chemosphere [Internet]. 2017 Jul [cited 2020 Apr 24];178:513-33. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S004565351730437X>
  45. Henderson G. A comparison of the effects of chromate, molybdate and cadmium oxide on respiration in the yeast *Saccharomyces cerevisiae*. Biol Met [Internet]. 1989 [cited 2020 Apr 12];2(2):83-8. Available from: <http://link.springer.com/10.1007/BF01129205>
  46. Cozza D, Torelli A, Veltri A, Ferrari M, Marieschi M, Cozza R. Ultrastructural features, chromium content and *in situ* immunodetection of 5-methyl-cytosine following Cr (VI) treatment in two strains of *Scenedesmus acutus* M. (Chlorophyceae) with different chromium sensitivity. Eur J Phycol [Internet]. 2016 Jul 2 [cited 2021 Sep 10];51(3):294-306. Available from: <https://www.tandfonline.com/doi/full/10.1080/09670262.2016.1157902>
  47. Hörcsik ZT, Kovács L, Láposi R, Mészáros I, Lakatos G, Garab G. Effect of chromium on photosystem 2 in the unicellular green alga, *Chlorella pyrenoidosa*. Photosynthetica [Internet]. 2007 Mar [cited 2020 May 25];45(1):65-9. Available from: <http://link.springer.com/10.1007/s11099-007-0010-8>
  48. Jin M, Xiao X, Qin L, Geng W, Gao Y, Li L, et al. Physiological and morphological responses and tolerance mechanisms of *Isochrysis galbana* to Cr(VI) stress. Bioresour Technol [Internet]. 2020 Apr [cited 2021 Sep 10];302:122860. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852420301292>
  49. Toranzo R, Ferraro G, Beligni MV, Perez GL, Castiglioni D, Pasquevich D, et al. Natural and acquired mechanisms of tolerance to chromium in a *Scenedesmus dimorphus* strain. Algal Res [Internet]. 2020 Dec [cited 2021 Sep 10];52:102100. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2211926420309681>
  50. Ünal D, Işık NO, Sukatar A. Effects of Chromium VI stress on green alga *Ulva lactuca* (L.). 2008;6.
  51. Volland S, Lütz C, Michalke B, Lütz-Meindl U. Intracellular chromium localization and cell physiological response in the unicellular alga *Micrasterias*. Aquat Toxicol [Internet]. 2012 Mar [cited 2021 Sep 10];109:59-69. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0166445X11003298>
  52. Saha B, Orvig C. Biosorbents for hexavalent chromium elimination from industrial and municipal effluents. Coord Chem Rev [Internet]. 2010 Dec [cited 2020 Apr 12];254(23-24):2959-72. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0010854510001591>
  53. Altun T, Parlayıcı Ş, Pehlivan E. Hexavalent chromium removal using agricultural waste "rye husk." Desalination Water Treat [Internet]. 2016 Aug 14 [cited 2020 Apr 13];57(38):17748-56. Available from: <http://www.tandfonline.com/doi/full/10.1080/19443994.2015.1085914>
  54. Dong C, Ji J, Shen B, Xing M, Zhang J. Enhancement of H<sub>2</sub>O<sub>2</sub> Decomposition by the Co-catalytic Effect of WS<sub>2</sub> on the Fenton Reaction for the Synchronous Reduction of Cr(VI) and Remediation of Phenol. Environ Sci Technol [Internet]. 2018 Oct 2 [cited 2021 Sep 10];52(19):11297-308. Available from: <https://pubs.acs.org/doi/10.1021/acs.est.8b02403>
  55. Jacobs J, L. Hardison R, Rouse V. J. In-situ remediation of heavy metals using sulfur-based treatment technologies. 2001;4.
  56. Ma Y, Li F, Jiang Y, Yang W, Lv L, Xue H, et al. Remediation of Cr(VI)-Contaminated Soil Using the Acidified Hydrazine Hydrate. Bull Environ Contam Toxicol [Internet]. 2016 Sep [cited 2021 Sep 10];97(3):392-4. Available from: <http://link.springer.com/10.1007/s00128-016-1862-z>
  57. Kabay N, Arda M, Saha B, Streat M. Removal of Cr(VI) by solvent impregnated resins (SIR) containing aliquat 336. React Funct Polym [Internet]. 2003 Jan [cited 2020 Apr 15];54(1-3):103-15. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1381514802001864>
  58. Jin W, Du H, Zheng S, Zhang Y. Electrochemical processes for the environmental remediation of toxic Cr(VI): A review. Electrochimica Acta [Internet]. 2016 Feb [cited 2021 Sep 11];191:1044-55. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0013468616301335>
  59. Li D, Ji G, Hu J, Hu S, Yuan X. Remediation strategy and electrochemistry flushing & reduction technology for real Cr(VI)-contaminated soils. Chem Eng J [Internet]. 2018 Feb [cited 2021 Sep 11];334:1281-8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1385894717319861>
  60. Belder C, Bedia J, Gómez-Avilés A, Peñas-Garzón M, Rodríguez JJ. Semiconductor Photocatalysis for Water Purification. In: Nanoscale Materials in Water Purification [Internet]. Elsevier; 2019 [cited 2021 Aug 2]. p. 581-651. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780128139264000288>



61. Izzudin NM, Jalil AA, Aziz FFA, Azami MS, Ali MW, Hassan NS, et al. Simultaneous remediation of hexavalent chromium and organic pollutants in wastewater using period 4 transition metal oxide-based photocatalysts: a review. *Environ Chem Lett* [Internet]. 2021 Jul 21 [cited 2021 Sep 11]; Available from: <https://link.springer.com/10.1007/s10311-021-01272-1>
62. Idris A, Hassan N, Mohd Ismail NS, Misran E, Yusof NM, Ngomsik A-F, et al. Photocatalytic magnetic separable beads for chromium (VI) reduction. *Water Res* [Internet]. 2010 Mar [cited 2020 Apr 15];44(6):1683-8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0043135409007684>
63. Xia S, Song Z, Jeyakumar P, Shaheen SM, Rinklebe J, Ok YS, et al. A critical review on bioremediation technologies for Cr(VI)-contaminated soils and wastewater. *Crit Rev Environ Sci Technol* [Internet]. 2019 Jun 18 [cited 2021 Jul 12];49(12):1027-78. Available from: <https://www.tandfonline.com/doi/full/10.1080/10643389.2018.1564526>
64. Zhang D, Xu Y, Li X, Wang L, He X, Ma Y, et al. The Immobilization Effect of Natural Mineral Materials on Cr(VI) Remediation in Water and Soil. *Int J Environ Res Public Health* [Internet]. 2020 Apr 20 [cited 2021 Sep 11];17(8):2832. Available from: <https://www.mdpi.com/1660-4601/17/8/2832>
65. Dehghani MH, Sanaei D, Ali I, Bhatnagar A. Removal of chromium(VI) from aqueous solution using treated waste newspaper as a low-cost adsorbent: Kinetic modeling and isotherm studies. *J Mol Liq* [Internet]. 2016 Mar [cited 2020 Apr 12];215:671-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0167732215313039>
66. Kumar S, Meikap BC. Removal of Chromium(VI) from waste water by using adsorbent prepared from green coconut shell. *Desalination Water Treat* [Internet]. 2014 May 12 [cited 2020 Apr 9];52(16-18):3122-32. Available from: <http://www.tandfonline.com/doi/abs/10.1080/19443994.2013.801796>
67. Owlad M, Aroua MK, Daud WAW, Baroutian S. Removal of Hexavalent Chromium-Contaminated Water and Wastewater: A Review. *Water Air Soil Pollut* [Internet]. 2009 Jun [cited 2020 Apr 9];200(1-4):59-77. Available from: <http://link.springer.com/10.1007/s11270-008-9893-7>
68. Ramirez Losada VA, Bonilla EP, Carvajal Pinilla LA, Serrezuela RR. Removal of chromium in wastewater from tanneries applying bioremediation with algae, orange peels and citrus pectin. *Contemp Eng Sci* [Internet]. 2018 [cited 2020 Apr 9];11(9):433-49. Available from: <http://www.m-hikari.com/ces/ces2018/ces9-12-2018/8235.html>
69. Zakaria ZA, Zakaria Z, Surif S, Ahmad WA. Biological detoxification of Cr(VI) using wood-husk immobilized *Acinetobacter haemolyticus*. *J Hazard Mater* [Internet]. 2007 Sep [cited 2020 Jun 4];148(1-2):164-71. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389407002439>
70. Choudhury P, Mondal P, Majumdar S, Saha S, Sahoo GC. Preparation of ceramic ultrafiltration membrane using green synthesized CuO nanoparticles for chromium (VI) removal and optimization by response surface methodology. *J Clean Prod* [Internet]. 2018 Dec [cited 2021 Sep 11];203:511-20. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0959652618326507>
71. Kumar A, Thakur A, Panesar PS. Extraction of hexavalent chromium by environmentally benign green emulsion liquid membrane using tridodecylamine as an extractant. *J Ind Eng Chem* [Internet]. 2019 Feb [cited 2021 Sep 11];70:394-401. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1226086X18308219>
72. Sharma SK, Petrusevski B, Amy G. Chromium removal from water: a review. *J Water Supply Res Technol-Aqua* [Internet]. 2008 Dec [cited 2020 Apr 15];57(8):541-53. Available from: <https://iwaponline.com/aqua/article/57/8/541/31147/Chromium-removal-from-water-a-review>
73. Hafiane A, Lemordant D, Dhahbi M. Removal of hexavalent chromium by nanofiltration. *Desalination* [Internet]. 2000 Nov [cited 2020 Apr 16];130(3):305-12. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0011916400000941>
74. Peng H, Guo J. Removal of chromium from wastewater by membrane filtration, chemical precipitation, ion exchange, adsorption electrocoagulation, electrochemical reduction, electrodialysis, electrodeionization, photocatalysis and nanotechnology: a review. *Environ Chem Lett* [Internet]. 2020 Nov [cited 2021 Sep 11];18(6):2055-68. Available from: <https://link.springer.com/10.1007/s10311-020-01058-x>
75. Sapari N, Idris A, Hamid NHAb. Total removal of heavy metal from mixed plating rinse wastewater. *Desalination* [Internet]. 1996 Aug [cited 2020 Apr 15];106(1-3):419-22. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0011916496001397>
76. Dangi AK, Sharma B, Hill RT, Shukla P. Bioremediation through microbes: systems biology and metabolic engineering approach. *Crit Rev Biotechnol* [Internet]. 2019 Jan 2 [cited 2021 Sep 26];39(1):79-98. Available from: <https://www.tandfonline.com/doi/full/10.1080/07388551.2018.1500997>
77. Verma S, Kuila A. Bioremediation of heavy metals by microbial process. *Environ Technol Innov* [Internet]. 2019 May [cited 2021 Sep 26];14:100369. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2352186418305911>
78. Rabani MS, Sharma R, Singh R, Gupta MK. Characterization and Identification of Naphthalene Degrading Bacteria Isolated from Petroleum Contaminated Sites and Their Possible Use in Bioremediation. *Polycycl Aromat Compd* [Internet]. 2020 May 6 [cited 2021 Sep 26];1-12. Available from: <https://www.tandfonline.com/doi/full/10.1080/10406638.2020.1759663>
79. Ontañón OM, Fernández M, Agostini E, González PS. Identification of the main mechanisms involved in the tolerance and bioremediation of Cr(VI) by *Bacillus* sp. SFC 500-1E. *Environ Sci Pollut Res* [Internet]. 2018 Jun [cited 2021 Jun 5];25(16):16111-20. Available from: <http://link.springer.com/10.1007/s11356-018-1764-1>
80. Diaconu M, Pavel LV, Hlihor R-M, Rosca M, Fertu DI, Lenz M, et al. Characterization of heavy metal toxicity in some plants and microorganisms – A preliminary approach for environmental bioremediation. *New Biotechnol* [Internet]. 2020 May [cited 2021 Sep 26];56:130-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1871678419300603>
81. Chen H, Wang Q. Microalgae-based nitrogen bioremediation. *Algal Res* [Internet]. 2020 Mar [cited 2021 Sep 26];46:101775. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2211926419311233>
82. Boopathy R. Factors limiting bioremediation technologies. *Bioresour Technol* [Internet]. 2000 Aug [cited 2020 Apr 19];74(1):63-7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852499001443>
83. Brar SK, Verma M, Surampalli RY, Misra K, Tyagi RD, Meunier N, et al. Bioremediation of Hazardous Wastes – A Review. *Pract Period Hazard Toxic Radioact Waste Manag* [Internet]. 2006 Apr [cited 2020 May 6];10(2):59-72. Available from: <http://ascelibrary.org/doi/10.1061/%28ASCE%291090-025X%282006%2910%3A2%2859%29>
84. Salunkhe PB, Dhakephalkar PK, Paknikar KM. Bioremediation of hexavalent chromium in soil microcosms. *Biotechnol Lett* [Internet]. 1998 [cited 2020 May 5];20(8):749-51. Available from: <http://link.springer.com/10.1023/A:1005338820430>
85. Molokwane PE, Nkhalambayausi-Chirwa EM. Microbial culture dynamics and chromium (VI) removal in packed-column microcosm reactors. *Water Sci Technol* [Internet]. 2009 Jul 1 [cited 2020 May 5];60(2):381-8. Available from: <https://iwaponline.com/wst/article/60/2/381/17771/Microbial-culture-dynamics-and-chromium-VI-removal>
86. Pavithra KG, Kumar PS, Jaikumar V, Vardhan KH, SundarRajan P. Microalgae for biofuel production and removal of heavy metals: a review. *Environ Chem Lett* [Internet]. 2020 Nov [cited 2021 Sep



- 11];18(6):1905-23. Available from: <https://link.springer.com/10.1007/s10311-020-01046-1>
87. Rahman Z, Singh VP. Bioremediation of toxic heavy metals (THMs) contaminated sites: concepts, applications and challenges. *Environ Sci Pollut Res* [Internet]. 2020 Aug [cited 2021 Sep 11];27(22):27563-81. Available from: <https://link.springer.com/10.1007/s11356-020-08903-0>
88. Thatoi H, Das S, Mishra J, Rath BP, Das N. Bacterial chromate reductase, a potential enzyme for bioremediation of hexavalent chromium: A review. *J Environ Manage* [Internet]. 2014 Dec [cited 2020 Apr 9];146:383-99. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0301479714003545>
89. Mistry K, Desai C, Lal S, Patel K, Patel B. Hexavalent Chromium Reduction by *Staphylococcus* Sp. Isolated From Cr (VI) Contaminated Land Fill. 2010;17.
90. Ohtake H, Cervantes C, Silver S. Decreased chromate uptake in *Pseudomonas fluorescens* carrying a chromate resistance plasmid. *J Bacteriol* [Internet]. 1987 [cited 2020 Apr 28];169(8):3853-6. Available from: <https://JB.asm.org/content/169/8/3853>
91. Suresh G, Balasubramanian B, Ravichandran N, Ramesh B, Kamyab H, Velmurugan P, et al. Bioremediation of hexavalent chromium-contaminated wastewater by *Bacillus thuringiensis* and *Staphylococcus capitis* isolated from tannery sediment. *Biomass Convers Biorefinery* [Internet]. 2021 Apr [cited 2021 Sep 12];11(2):383-91. Available from: <http://link.springer.com/10.1007/s13399-020-01259-y>
92. Vatsouria A, Vainshtein M, Kusch P, Wiessner A, D K, Kaestner M. Anaerobic co-reduction of chromate and nitrate by bacterial cultures of *Staphylococcus epidermidis* L-02. *J Ind Microbiol Biotechnol* [Internet]. 2005 Sep [cited 2021 Jun 9];32(9):409-14. Available from: <https://academic.oup.com/jimb/article/32/9/409-414/5992807>
93. Bopp LH, Ehrlich HL. Chromate resistance and reduction in *Pseudomonas fluorescens* strain LB300. *Arch Microbiol* [Internet]. 1988 Sep [cited 2020 Apr 9];150(5):426-31. Available from: <http://link.springer.com/10.1007/BF00422281>
94. Campos J, Martinez-Pacheco M, Cervantes C. Hexavalent-chromium reduction by a chromate-resistant *Bacillus* sp. strain. *Antonie Van Leeuwenhoek* [Internet]. 1995 Oct [cited 2020 Apr 9];68(3):203-8. Available from: <http://link.springer.com/10.1007/BF00871816>
95. Opperman DJ, van Heerden E. Aerobic Cr(VI) reduction by *Thermus scotoductus* strain SA-01: Cr(VI) reduction by *Thermus scotoductus*. *J Appl Microbiol* [Internet]. 2007 Nov [cited 2020 Apr 21];103(5):1907-13. Available from: <http://doi.wiley.com/10.1111/j.1365-2672.2007.03429.x>
96. Wang Y. Factors affecting hexavalent chromium reduction in pure cultures of bacteria. *Water Res* [Internet]. 1995 Nov [cited 2020 Apr 28];29(11):2467-74. Available from: <https://linkinghub.elsevier.com/retrieve/pii/004313549500093Z>
97. Narayani M, Shetty KV. Chromium-Resistant Bacteria and Their Environmental Condition for Hexavalent Chromium Removal: A Review. *Crit Rev Environ Sci Technol* [Internet]. 2013 Jan [cited 2020 Apr 17];43(9):955-1009. Available from: <http://www.tandfonline.com/doi/abs/10.1080/10643389.2011.627022>
98. Tamindžija D, Chromikova Z, Spaić A, Barak I, Bernier-Latmani R, Radnović D. Chromate tolerance and removal of bacterial strains isolated from uncontaminated and chromium-polluted environments. *World J Microbiol Biotechnol* [Internet]. 2019 Apr [cited 2021 Sep 12];35(4):56. Available from: <http://link.springer.com/10.1007/s11274-019-2638-5>
99. Srivastava S, Thakur IS. Isolation and process parameter optimization of *Aspergillus* sp. for removal of chromium from tannery effluent. *Bioresour Technol* [Internet]. 2006 Jul [cited 2020 Apr 22];97(10):1167-73. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852405002725>
100. Martorell MM, Fernández PM, Fariña JI, Figueroa LIC. Cr(VI) reduction by cell-free extracts of *Pichia jadinii* and *Pichia anomala* isolated from textile-dye factory effluents. *Int Biodeterior Biodegrad* [Internet]. 2012 Jul [cited 2020 Apr 16];71:80-5. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830512000935>
101. Morales-Barrera L, Guillén-Jiménez F de M, Ortiz-Moreno A, Villegas-Garrido TL, Sandoval-Cabrera A, Hernández-Rodríguez CH, et al. Isolation, identification and characterization of a *Hypocrea tawa* strain with high Cr(VI) reduction potential. *Biochem Eng J* [Internet]. 2008 Jun [cited 2020 Apr 16];40(2):284-92. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1369703X0800003X>
102. Morales-Barrera L, Cristiani-Urbina E. Hexavalent Chromium Removal by a *Trichoderma inhamatum* Fungal Strain Isolated from Tannery Effluent. *Water Air Soil Pollut* [Internet]. 2007 Nov 29 [cited 2020 Apr 16];187(1-4):327-36. Available from: <http://link.springer.com/10.1007/s11270-007-9520-z>
103. Vankar PS, Bajpai D. Phyto-remediation of chrome-VI of tannery effluent by *Trichoderma* species. *Desalination* [Internet]. 2008 Mar [cited 2020 Apr 16];222(1-3):255-62. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S001191640700776X>
104. Mahmoud MS, Mohamed SA. Calcium alginate as an eco-friendly supporting material for Baker's yeast strain in chromium bioremediation. *HBRC J* [Internet]. 2017 Dec [cited 2020 Apr 17];13(3):245-54. Available from: <https://www.tandfonline.com/doi/full/10.1016/j.hbrj.2015.06.003>
105. Campaña-Pérez JF, Portero Barahona P, Martín-Ramos P, Carvajal Barriga EJ. Ecuadorian yeast species as microbial particles for Cr(VI) biosorption. *Environ Sci Pollut Res* [Internet]. 2019 Sep [cited 2021 Sep 16];26(27):28162-72. Available from: <http://link.springer.com/10.1007/s11356-019-06035-8>
106. Adam S, P SK, P S, S DK, P P. Bioremediation of Tannery Wastewater Using Immobilized Marine Microalga *Tetraselmis* sp.: Experimental Studies and Pseudo-Second Order Kinetics. *J Mar Biol Oceanogr* [Internet]. 2015 [cited 2020 Apr 9];04(01). Available from: [https://www.scitechnol.com/peer-review/bioremediation-of-tannery-wastewater-using-immobilized-marine-microalga-tetraselmis-sp-experimental-studies-and-pseudo-second-order-Uoxr.php?article\\_id=3505](https://www.scitechnol.com/peer-review/bioremediation-of-tannery-wastewater-using-immobilized-marine-microalga-tetraselmis-sp-experimental-studies-and-pseudo-second-order-Uoxr.php?article_id=3505)
107. Gupta VK, Shrivastava AK, Jain N. Biosorption of Chromium(VI) From Aqueous solutions by green algae *spirogyra* species. *Water Res* [Internet]. 2001 Dec [cited 2020 Apr 17];35(17):4079-85. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0043135401001385>
108. Lee D-C, Park C-J, Yang J-E, Jeong Y-H, Rhee H-I. Screening of hexavalent chromium biosorbent from marine algae. *Appl Microbiol Biotechnol* [Internet]. 2000 Sep 15 [cited 2020 Apr 17];54(3):445-8. Available from: <http://link.springer.com/10.1007/s0025300000387>
109. Yewalkar SN, Dhumal Kondiram N, Sainis JK. Chromium(VI)-reducing *Chlorella* spp. isolated from disposal sites of paper-pulp and electroplating industry. *J Appl Phycol* [Internet]. 2007 Aug 28 [cited 2020 May 28];19(5):459-65. Available from: <http://link.springer.com/10.1007/s10811-007-9153-z>
110. El-Sikaily A, Nemr AE, Khaled A, Abdelwehab O. Removal of toxic chromium from wastewater using green alga *Ulva lactuca* and its activated carbon. *J Hazard Mater* [Internet]. 2007 Sep [cited 2020 Apr 17];148(1-2):216-28. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389407002506>
111. Aksu Z, Açikel Ü, Kutsal T. Investigation of Simultaneous Biosorption of Copper(II) and Chromium(VI) on Dried *Chlorella Vulgaris* from Binary Metal Mixtures: Application of Multicomponent Adsorption Isotherms. *Sep Sci Technol* [Internet]. 1999 Feb 22 [cited 2020 May 28];34(3):501-24. Available from: <http://www.tandfonline.com/doi/abs/10.1081/SS-100100663>
112. Athira K, Sathish A, Nithya K, Guhananthan A. Corn cob immobilised *Chlorella sorokiniana* for the sequestration of chromium ions from aqueous solution. *Mater Today Proc* [Internet]. 2020 Apr [cited 2020 May 28];S2214785320319313. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2214785320319313>



113. Deng L, Wang H, Deng N. Photoreduction of chromium(VI) in the presence of algae, *Chlorella vulgaris*. J Hazard Mater [Internet]. 2006 Nov 16 [cited 2020 Apr 15];138(2):288–92. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389406006297>
114. Han X, Wong YS, Wong MH, Tam NFY. Biosorption and bioreduction of Cr(VI) by a microalgal isolate, *Chlorella miniata*. J Hazard Mater [Internet]. 2007 Jul [cited 2020 Apr 9];146(1–2):65–72. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S030438940601418X>
115. Costa IGF, Terra NM, Cardoso VL, Batista FRX, Reis MHM. Photoreduction of chromium(VI) in microstructured ceramic hollow fibers impregnated with titanium dioxide and coated with green algae *Chlorella vulgaris*. J Hazard Mater [Internet]. 2019 Nov [cited 2021 Sep 16];379:120837. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389419307903>
116. Dittert IM, de Lima Brandão H, Pina F, da Silva EAB, de Souza SMAGU, de Souza AAU, et al. Integrated reduction/oxidation reactions and sorption processes for Cr(VI) removal from aqueous solutions using *Laminaria digitata* macro-algae. Chem Eng J [Internet]. 2014 Feb [cited 2021 Aug 3];237:443–54. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1385894713013697>
117. Adki VS, Jadhav JP, Bapat VA. Nopalea cochenillifera, a potential chromium (VI) hyperaccumulator plant. Environ Sci Pollut Res [Internet]. 2013 Feb [cited 2020 Apr 24];20(2):1173–80. Available from: <http://link.springer.com/10.1007/s11356-012-1125-4>
118. Ranieri E, Gikas P. Effects of Plants for Reduction and Removal of Hexavalent Chromium from a Contaminated Soil. Water Air Soil Pollut [Internet]. 2014 Jun [cited 2020 Apr 24];225(6):1981. Available from: <http://link.springer.com/10.1007/s11270-014-1981-2>
119. Shanker A, Djanaguiraman M, Sudhagar R, Chandrashekar C, Pathmanabhan G. Differential antioxidative response of ascorbate glutathione pathway enzymes and metabolites to chromium speciation stress in green gram ( (*L.*) *R. Wilczek*. cv CO 4) roots. Plant Sci [Internet]. 2004 Apr [cited 2020 Apr 29];166(4):1035–43. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0168945203005247>
120. Ma H-L, Zhang Y, Hu Q-H, Yan D, Yu Z-Z, Zhai M. Chemical reduction and removal of Cr(vi) from acidic aqueous solution by ethylenediamine-reduced graphene oxide. J Mater Chem [Internet]. 2012 [cited 2020 Apr 15];22(13):5914. Available from: <http://xlink.rsc.org/?DOI=c2jm00145d>
121. Graham MC, Farmer JG, Anderson P, Paterson E, Hillier S, Lumsdon DG, et al. Calcium polysulfide remediation of hexavalent chromium contamination from chromite ore processing residue. Sci Total Environ [Internet]. 2006 Jul [cited 2020 May 21];364(1–3):32–44. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0048969705008387>
122. Park D, Yun Y-S, Park JM. Studies on hexavalent chromium biosorption by chemically-treated biomass of *Ecklonia* sp. Chemosphere [Internet]. 2005 Sep [cited 2020 May 21];60(10):1356–64. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653505002766>
123. Barrera-Díaz C, Palomar-Pardavé M, Romero-Romo M, Martínez S. Chemical and electrochemical considerations on the removal process of hexavalent chromium from aqueous media. J Appl Electrochem [Internet]. 2003 [cited 2020 May 21];33(11):61–71. Available from: <http://link.springer.com/10.1023/A:1022983919644>
124. Wang Y, Fang Z, Liang B, Tsang EP. Remediation of hexavalent chromium contaminated soil by stabilized nanoscale zero-valent iron prepared from steel pickling waste liquor. Chem Eng J [Internet]. 2014 Jul [cited 2020 May 22];247:283–90. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S138589471400285X>
125. Qin G, McGuire MJ, Blute NK, Seidel C, Fong L. Hexavalent Chromium Removal by Reduction with Ferrous Sulfate, Coagulation, and Filtration: A Pilot-Scale Study. Environ Sci Technol [Internet]. 2005 Aug 1 [cited 2021 Aug 7];39(16):6321–7. Available from: <https://pubs.acs.org/doi/10.1021/es050486p>
126. Rahaman A, Hosen MdR, Hena MA, Naher UHB, Moniruzzaman M. A Study on removal of chromium from tannery effluent treatment of chrome tanning waste water using tannery solid waste. Int J Hum Cap Urban Manag [Internet]. 2016 Oct [cited 2020 Apr 9];1(4). Available from: <http://doi.org/10.22034/ijhcum.2016.04.001>
127. Krishna D, Sree RP. Artificial Neural Network and Response Surface Methodology Approach for Modeling and Optimization of Chromium (VI) Adsorption from Waste Water using Ragi Husk Powder. Indian Chem Eng [Internet]. 2013 Sep [cited 2020 Apr 9];55(3):200–22. Available from: <http://www.tandfonline.com/doi/abs/10.1080/00194506.2013.829257>
128. Garg UK, Kaur MP, Garg VK, Sud D. Removal of hexavalent chromium from aqueous solution by agricultural waste biomass. J Hazard Mater [Internet]. 2007 Feb [cited 2020 Apr 12];140(1–2):60–8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S030438940600690X>
129. Saha R, Saha B. Removal of hexavalent chromium from contaminated water by adsorption using mango leaves (*Mangifera indica*). Desalination Water Treat [Internet]. 2014 Mar 21 [cited 2020 Apr 19];52(10–12):1928–36. Available from: <http://www.tandfonline.com/doi/abs/10.1080/19443994.2013.804458>
130. Shi T, Wang Z, Liu Y, Jia S, Changming D. Removal of hexavalent chromium from aqueous solutions by D301, D314 and D354 anion-exchange resins. J Hazard Mater [Internet]. 2009 Jan [cited 2021 Jul 19];161(2–3):900–6. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389408005748>
131. Chen J-H, Hsu K-C, Chang Y-M. Surface Modification of Hydrophobic Resin with Tricaprylmethylammonium Chloride for the Removal of Trace Hexavalent Chromium. Ind Eng Chem Res [Internet]. 2013 Aug 21 [cited 2021 Jul 19];52(33):11685–94. Available from: <https://pubs.acs.org/doi/10.1021/ie401233r>
132. Bansal M, Garg U, Singh D, Garg VK. Removal of Cr(VI) From Aqueous Solutions Using Pre-Consumer Processing Agricultural Waste: A Case Study of Rice Husk. J Hazard Mater. 2008 Jun 1;162:312–20.
133. Setshedi KZ, Bhaumik M, Songwane S, Onyango MS, Maity A. Exfoliated polypyrrole-organically modified montmorillonite clay nanocomposite as a potential adsorbent for Cr(VI) removal. Chem Eng J [Internet]. 2013 Apr [cited 2021 Aug 7];222:186–97. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1385894713002271>
134. Wang X, Lu J, Cao B, Liu X, Lin Z, Yang C, et al. Facile synthesis of recycling Fe<sub>3</sub>O<sub>4</sub>/graphene adsorbents with potassium humate for Cr(VI) removal. Colloids Surf Physicochem Eng Asp [Internet]. 2019 Jan [cited 2021 Aug 7];560:384–92. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0927775718310288>
135. Hokkanen S, Bhatnagar A, Repo E, Lou S, Sillanpää M. Calcium hydroxyapatite microfibrillated cellulose composite as a potential adsorbent for the removal of Cr(VI) from aqueous solution. Chem Eng J [Internet]. 2016 Jan [cited 2021 Aug 7];283:445–52. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1385894715010013>
136. Padmavathy KS, Madhu G, Haseena PV. A study on Effects of pH, Adsorbent Dosage, Time, Initial Concentration and Adsorption Isotherm Study for the Removal of Hexavalent Chromium (Cr (VI)) from Wastewater by Magnetite Nanoparticles. Procedia Technol [Internet]. 2016 [cited 2021 Aug 7];24:585–94. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S221201731630216X>
137. Cherdchoo W, Nithetham S, Charoenpanich J. Removal of Cr(VI) from synthetic wastewater by adsorption onto coffee ground and mixed waste tea. Chemosphere [Internet]. 2019 Apr [cited 2021 Jun 3];221:758–67. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653519301109>
138. Dakiky M, Khamis M, Manassra A, Mer'eb M. Selective adsorption of chromium(VI) in industrial wastewater using low-cost



- abundantly available adsorbents. *Adv Environ Res* [Internet]. 2002 Oct [cited 2021 Aug 7];6(4):533-40. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S109301910100079X>
139. Mdalose L, Balogun M, Setshedi K, Chimuka L, Chetty A. Performance evaluation of polypyrrole-montmorillonite clay composite as a re-usable adsorbent for Cr(VI) remediation. *Polym Bull* [Internet]. 2021 Aug [cited 2021 Sep 11];78(8):4685-97. Available from: <https://link.springer.com/10.1007/s00289-020-03338-6>
  140. Tariq MA, Nadeem M, Iqbal MM, Imran M, Siddique MH, Iqbal Z, et al. Effective sequestration of Cr (VI) from wastewater using nanocomposite of ZnO with cotton stalks biochar: modeling, kinetics, and reusability. *Environ Sci Pollut Res* [Internet]. 2020 Sep [cited 2021 Sep 11];27(27):33821-34. Available from: <https://link.springer.com/10.1007/s11356-020-09481-x>
  141. Saxena D, Levin R, Firer MA. Removal of chromate from industrial effluent by a new isolate of *Staphylococcus cohnii*. *Water Sci Technol* [Internet]. 2000 Jul 1 [cited 2020 Apr 9];42(1-2):93-8. Available from: <https://iwaponline.com/wst/article/42/1-2/93/9910/Removal-of-chromate-from-industrial-effluent-by-a>
  142. Yao Y, Hu L, Li S, Zeng Q, Zhong H, He Z. Exploration on the bioreduction mechanisms of Cr(VI) and Hg(II) by a newly isolated bacterial strain *Pseudomonas umsongensis* CY-1. *Ecotoxicol Environ Saf*. 2020 Sep 1;201:110850.
  143. Tan H, Wang C, Zeng G, Luo Y, Li H, Xu H. Bioreduction and biosorption of Cr(VI) by a novel *Bacillus* sp. CRB-B1 strain. *J Hazard Mater* [Internet]. 2020 Mar [cited 2021 Aug 7];386:121628. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389419315821>
  144. Huang X-N, Min D, Liu D-F, Cheng L, Qian C, Li W-W, et al. Formation mechanism of organo-chromium (III) complexes from bioreduction of chromium (VI) by *Aeromonas hydrophila*. *Environ Int* [Internet]. 2019 Aug [cited 2021 Aug 7];129:86-94. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0160412019308232>
  145. Sun Y, Lan J, Du Y, Guo L, Du D, Chen S, et al. Chromium(VI) bioreduction and removal by *Enterobacter* sp. SL grown with waste molasses as carbon source: Impact of operational conditions. *Bioresour Technol* [Internet]. 2020 Apr [cited 2021 Sep 12];302:121974. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852419312040>
  146. Princy S, Sathish SS, Cibichakravarthy B, Prabakaran SR. Hexavalent chromium reduction by *Morganella morganii* (1Ab1) isolated from tannery effluent contaminated sites of Tamil Nadu, India. *Biocatal Agric Biotechnol* [Internet]. 2020 Jan [cited 2021 Sep 12];23:101469. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1878818119316846>
  147. Gao J, Wu S, Liu Y, Wu S, Jiang C, Li X, et al. Characterization and transcriptomic analysis of a highly Cr(VI)-resistant and -reductive plant-growth-promoting rhizobacterium *Stenotrophomonas rhizophila* DSM14405T. *Environ Pollut* [Internet]. 2020 Aug [cited 2021 Sep 12];263:114622. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0269749119355216>
  148. Tirry N, Tahri Joutey N, Sayel H, Kouchou A, Bahafid W, Asri M, et al. Screening of plant growth promoting traits in heavy metals resistant bacteria: Prospects in phytoremediation. *J Genet Eng Biotechnol* [Internet]. 2018 Dec [cited 2021 Sep 12];16(2):613-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1687157X18300635>
  149. Elmeihy R, Shi X-C, Tremblay P-L, Zhang T. Fast removal of toxic hexavalent chromium from an aqueous solution by high-density *Geobacter sulfurreducens*. *Chemosphere* [Internet]. 2021 Jan [cited 2021 Sep 12];263:128281. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653520324760>
  150. An Q, Deng S, Xu J, Nan H, Li Z, Song J-L. Simultaneous reduction of nitrate and Cr(VI) by *Pseudomonas aeruginosa* strain G12 in wastewater. *Ecotoxicol Environ Saf* [Internet]. 2020 Mar [cited 2021 Jul 27];191:110001. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0147651319313326>
  151. Cheng J, Gao J, Zhang J, Yuan W, Yan S, Zhou J, et al. Optimization of Hexavalent Chromium Biosorption by *Shewanella putrefaciens* Using the Box-Behnken Design. *Water Air Soil Pollut* [Internet]. 2021 Mar [cited 2021 Sep 12];232(3):92. Available from: <http://link.springer.com/10.1007/s11270-020-04947-7>
  152. Ma L, Xu J, Chen N, Li M, Feng C. Microbial reduction fate of chromium (Cr) in aqueous solution by mixed bacterial consortium. *Ecotoxicol Environ Saf* [Internet]. 2019 Apr [cited 2021 Aug 7];170:763-70. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0147651318313319>
  153. Prabhakaran D, Subramanian S. Studies on the Bioremediation of Chromium from Aqueous Solutions Using *C. paurometabolum*. *Trans Indian Inst Met*. 2016 Nov 22;70.
  154. Karthik C, Vijayan SR, Pugazhendhi A, Kumar G, Arulselvi P. Biosorption and biotransformation of Cr(VI) by novel Cellulosimicrobium funkei strain AR6. *J Taiwan Inst Chem Eng*. 2017 Jan 26;17:0-0.
  155. Sathishkumar K, Murugan K, Benelli G, Higuchi A, Rajasekar A. Bioreduction of hexavalent chromium by *Pseudomonas stutzeri* L1 and *Acinetobacter baumannii* L2. *Ann Microbiol* [Internet]. 2017 Jan [cited 2021 Aug 7];67(1):91-8. Available from: <http://link.springer.com/10.1007/s13213-016-1240-4>
  156. Chen C-Y, Cheng C-Y, Chen C-K, Hsieh M-C, Lin S-T, Ho K-Y, et al. Hexavalent chromium removal and bioelectricity generation by *Ochrobactrum* sp. YC211 under different oxygen conditions. *J Environ Sci Health Part A* [Internet]. 2016 May 11 [cited 2021 Aug 7];51(6):502-8. Available from: <http://www.tandfonline.com/doi/full/10.1080/10934529.2015.1128731>
  157. Sharma S, Adholeya A. Detoxification and accumulation of chromium from tannery effluent and spent chrome effluent by *Paecilomyces lilacinus* fungi. *Int Biodeterior Biodegrad* [Internet]. 2011 Mar [cited 2021 Aug 7];65(2):309-17. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830510002118>
  158. Pal S, Y V. Bioremediation of Chromium from Fortified Solutions by *Phanerochaete chrysosporium* (MTCC 787). *J Bioremediation Biodegrad* [Internet]. 2011 [cited 2021 Aug 7];02(05). Available from: <https://www.omicsonline.org/bioremediation-of-chromium-from-fortified-solutions-by-phanerochaete-chrysosporium-mtcc-787-2155-6199.1000127.php?aid=2103>
  159. Carol D, Kingsley S, Vincent S. Hexavalent chromium removal from aqueous solutions by *Pleurotus ostreatus* spent biomass. *Int J Eng Sci*. 2012 Jan 1;4:7-22.
  160. Chen R, Cheng Y, Wang P, Liu Z, Wang Y, Wang Y. High efficient removal and mineralization of Cr(VI) from water by functionalized magnetic fungus nanocomposites. *J Cent South Univ* [Internet]. 2020 May [cited 2021 Sep 16];27(5):1503-14. Available from: <https://link.springer.com/10.1007/s11771-020-4386-y>
  161. Husien Sh, Labena A, El-Beley EF, Mahmoud HM, Hamouda AS. Absorption of hexavalent chromium by green micro algae *Chlorella sorokiniana*: live planktonic cells. *Water Pract Technol* [Internet]. 2019 Sep 1 [cited 2020 Apr 17];14(3):515-29. Available from: <https://iwaponline.com/wpt/article/14/3/515/67497/Absorption-of-hexavalent-chromium-by-green-micro>
  162. Deng L, Zhang Y, Qin J, Wang X, Zhu X. Biosorption of Cr(VI) from aqueous solutions by nonliving green algae *Cladophora albida*. *Miner Eng* [Internet]. 2009 Mar [cited 2020 Apr 9];22(4):372-7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0892687508002525>
  163. Kumaraguru K, Saravanan P, Rajesh kannan R, Saravanan V. A systematic analysis of hexavalent chromium adsorption and elimination from aqueous environment using brown marine algae (*Turbinaria ornata*). *Biomass Convers Biorefinery* [Internet]. 2021 Aug 6 [cited 2021 Sep 16]; Available from: <https://link.springer.com/10.1007/s13399-021-01795-1>
  164. Mari S, Vrane J. Characteristics and significance of microbial biofilm formation. *Period Biol*. 2007;109(2):7.



165. Toyofuku M, Inaba T, Kiyokawa T, Obana N, Yawata Y, Nomura N. Environmental factors that shape biofilm formation. *Biosci Biotechnol Biochem* [Internet]. 2016 Jan 2 [cited 2019 Feb 5];80(1):7-12. Available from: <https://www.tandfonline.com/doi/full/10.1080/09168451.2015.1058701>
166. Evans LV. *Biofilms recent advances in their study and control*. Amsterdam: Taylor & Francis e-Library; 2004.
167. O'Toole G, Kaplan HB, Kolter R. Biofilm Formation as Microbial Development. *Annu Rev Microbiol* [Internet]. 2000 Oct [cited 2020 Apr 9];54(1):49-79. Available from: <http://www.annualreviews.org/doi/10.1146/annurev.micro.54.1.49>
168. Edwards SJ, Kjellerup BV. Applications of biofilms in bioremediation and biotransformation of persistent organic pollutants, pharmaceuticals/personal care products, and heavy metals. *Appl Microbiol Biotechnol* [Internet]. 2013 Dec [cited 2020 May 18];97(23):9909-21. Available from: <http://link.springer.com/10.1007/s00253-013-5216-z>
169. Das N, Basak LVG, Salam JA. Application of Biofilms on Remediation of Pollutants – An Overview. 2012;
170. Shukla SK, Mangwani N, Rao TS, Das S. Biofilm-Mediated Bioremediation of Polycyclic Aromatic Hydrocarbons. In: *Microbial Biodegradation and Bioremediation* [Internet]. Elsevier; 2014 [cited 2020 May 18]. p. 203-32. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B978012800021200008X>
171. Asri M, El Ghachtouli N, Elabed S, Ibsouda Koraichi S, Elabed A, Silva B, et al. *Wicherhamomyces anomalus* biofilm supported on wood husk for chromium wastewater treatment. *J Hazard Mater* [Internet]. 2018 Oct [cited 2020 Jun 1];359:554-62. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389418304060>
172. Herath HMLI, Rajapaksha AU, Vithanage M, Seneviratne G. Developed fungal-bacterial biofilms as a novel tool for bioremoval of hexavalent chromium from wastewater. *Chem Ecol* [Internet]. 2014 Jul 4 [cited 2020 Jun 1];30(5):418-27. Available from: <http://www.tandfonline.com/doi/abs/10.1080/02757540.2013.861828>
173. Smith WL, Gadd GM. Reduction and precipitation of chromate by mixed culture sulphate-reducing bacterial biofilms. *J Appl Microbiol* [Internet]. 2000 Jun [cited 2020 Jun 1];88(6):983-91. Available from: <http://doi.wiley.com/10.1046/j.1365-2672.2000.01066.x>
174. Yong P, Liu W, Zhang Z, Beaugard D, Johns ML, Macaskie LE. One step bioconversion of waste precious metals into *Serratia* biofilm-immobilized catalyst for Cr(VI) reduction. *Biotechnol Lett* [Internet]. 2015 Nov [cited 2020 Jun 1];37(11):2181-91. Available from: <http://link.springer.com/10.1007/s10529-015-1894-1>
175. Tripathi AG A. Bioremediation of toxic chromium from electroplating effluent by chromate-reducing *Pseudomonas aeruginosa* A2Chr in two bioreactors. *Appl Microbiol Biotechnol* [Internet]. 2002 Mar 1 [cited 2020 Apr 9];58(3):416-20. Available from: <http://link.springer.com/10.1007/s00253-001-0871-x>
176. Gabr RM, Gad-Elrab SMF, Abskharon RNN, Hassan SHA, Shoreit AAM. Biosorption of hexavalent chromium using biofilm of *E. coli* supported on granulated activated carbon. *World J Microbiol Biotechnol* [Internet]. 2009 Oct [cited 2020 Jun 2];25(10):1695-703. Available from: <http://link.springer.com/10.1007/s11274-009-0063-x>
177. Pan X, Liu Z, Chen Z, Cheng Y, Pan D, Shao J, et al. Investigation of Cr(VI) reduction and Cr(III) immobilization mechanism by planktonic cells and biofilms of *Bacillus subtilis* ATCC-6633. *Water Res* [Internet]. 2014 May [cited 2020 Jun 1];55:21-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0043135414001080>
178. Arica MY, Bayramoğlu G. Cr(VI) biosorption from aqueous solutions using free and immobilized biomass of *Lentinus sajor-caju*: preparation and kinetic characterization. *Colloids Surf Physicochem Eng Asp* [Internet]. 2005 Feb [cited 2021 Aug 6];253(1-3):203-11. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0927775704008908>
179. Carmona M, Silva M, Leite S, Vasco-Echeverri O, Ocampo-Lopez C. Packed bed redistribution system for Cr(III) and Cr(VI) biosorption by *Saccharomyces cerevisiae*. *J Taiwan Inst Chem Eng*. 2011 Jan 1;43.
180. Sanghi R, Srivastava A. Long-term chromate reduction by immobilized fungus in continuous column. *Chem Eng J*. 2010 Aug 1;162:122-6.
181. Ahmad A, Bhat AH, Buang A. Enhanced biosorption of transition metals by living *Chlorella vulgaris* immobilized in Ca-alginate beads. *Environ Technol* [Internet]. 2019 Jun 20 [cited 2021 Aug 6];40(14):1793-809. Available from: <https://www.tandfonline.com/doi/full/10.1080/09593330.2018.1430171>
182. Akhtar N, Iqbal M, Zafar SI, Iqbal J. Biosorption characteristics of unicellular green alga *Chlorella sorokiniana* immobilized in loofa sponge for removal of Cr(III). *J Environ Sci* [Internet]. 2008 Feb [cited 2021 Jul 16];20(2):231-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1001074208600364>
183. Baykal ÖZER T, Açıköz Erkaya İ, Udoh A, Özer T, Akbulut A, Bayramoglu G, et al. Biosorption of Cr(VI) by free and immobilized *Pediastrum boryanum* biomass: Equilibrium, kinetic, and thermodynamic studies. *Environ Sci Pollut Res Int*. 2012 Feb 29;19:2983-93.
184. petrovič A, Simonič M. Removal of heavy metal ions from drinking water by alginate-immobilised *Chlorella sorokiniana*. *Int J Environ Sci Technol*. 2016 May 18;13.
185. Wong Y-S, Tam NFY, editors. *Wastewater Treatment with Algae* [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg; 1998 [cited 2021 Aug 6]. Available from: <http://link.springer.com/10.1007/978-3-662-10863-5>
186. Abinandan S, Subashchandrabose SR, Venkateswarlu K, Megharaj M. Microalgae-bacteria biofilms: a sustainable synergistic approach in remediation of acid mine drainage. *Appl Microbiol Biotechnol* [Internet]. 2018 Feb [cited 2021 Sep 27];102(3):1131-44. Available from: <http://link.springer.com/10.1007/s00253-017-8693-7>
187. Qu W, Zhang C, Chen X, Ho S-H. New concept in swine wastewater treatment: development of a self-sustaining synergetic microalgae-bacteria symbiosis (ABS) system to achieve environmental sustainability. *J Hazard Mater* [Internet]. 2021 Sep [cited 2021 Sep 27];418:126264. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389421012280>
188. Sun J, Xu W, Cai B, Huang G, Zhang H, Zhang Y, et al. High-concentration nitrogen removal coupling with bioelectric power generation by a self-sustaining algal-bacterial biocathode photo-bioelectrochemical system under daily light/dark cycle. *Chemosphere* [Internet]. 2019 May [cited 2021 Sep 27];222:797-809. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0045653519302139>
189. Roestorff M, Chirwa E. Bacterial cr(vi) reduction with internal carbon recirculation using freshwater algae as primary producers. *Chem Eng Trans* [Internet]. 2018 May [cited 2020 Jun 4];64:457-62. Available from: <http://doi.org/10.33031/CET1864077>
190. Quintelas C, Fonseca B, Silva B, Figueiredo H, Tavares T. Treatment of chromium(VI) solutions in a pilot-scale bioreactor through a biofilm of *Arthrobacter viscosus* supported on GAC. *Bioresour Technol* [Internet]. 2009 Jan [cited 2020 Jun 3];100(1):220-6. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852408004422>
191. Morales DK, Ocampo W, Zambrano MM. Efficient removal of hexavalent chromium by a tolerant *Streptomyces* sp. affected by the toxic effect of metal exposure: Chromium tolerant *Streptomyces* sp. *J Appl Microbiol* [Internet]. 2007 Aug 30 [cited 2021 Jul 19];103(6):2704-12. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2672.2007.03510.x>
192. Córdoba A, Vargas P, Dussan J. Chromate reduction by *Arthrobacter* CR47 in biofilm packed bed reactors. *J Hazard Mater* [Internet]. 2008 Feb [cited 2020 Jun 3];151(1):274-9. Available from:

- <https://linkinghub.elsevier.com/retrieve/pii/S0304389407015555>
193. Sarioglu OF, Celebioglu A, Tekinay T, Uyar T. Bacteria-immobilized electrospun fibrous polymeric webs for hexavalent chromium remediation in water. *Int J Environ Sci Technol* [Internet]. 2016 Aug [cited 2020 Jun 3];13(8):2057–66. Available from: <http://link.springer.com/10.1007/s13762-016-1033-0>
  194. Dey S, Paul AK. Influence of metal ions on biofilm formation by *Arthrobacter* sp. SUK 1205 and evaluation of their Cr(VI) removal efficacy. *Int Biodeterior Biodegrad* [Internet]. 2018 Aug [cited 2020 Jun 3];132:122–31. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830517310466>
  195. Focardi S, Pepi M, Landi G, Gasperini S, Ruta M, Di Biasio P, et al. Hexavalent chromium reduction by whole cells and cell free extract of the moderate halophilic bacterial strain *Halomonas* sp. TA-04. *Int Biodeterior Biodegrad* [Internet]. 2012 Jan [cited 2020 Jun 3];66(1):63–70. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830511002307>
  196. Ravikumar KVG, Kumar D, Kumar G, Mrudula P, Natarajan C, Mukherjee A. Enhanced Cr(VI) Removal by Nanozerovalent Iron-Immobilized Alginate Beads in the Presence of a Biofilm in a Continuous-Flow Reactor. *Ind Eng Chem Res* [Internet]. 2016 May 25 [cited 2020 Jun 3];55(20):5973–82. Available from: <https://pubs.acs.org/doi/10.1021/acs.iecr.6b01006>
  197. Ait-Meddour A, Abbas N, Ouled-Haddar H, Sifour M, Bendjeddou K, Idoui T. Biofilm Formation by the Hexavalent Chromium Removing Strain *Streptococcus salivarius*: in Vitro Approach on Abiotic Surfaces. *Pollution* [Internet]. 2020 Apr [cited 2020 Jun 4];6(2). Available from: <http://doi.org/10.22059/poll.2020.288349.685>
  198. Chirwa EMN, Wang Y-T. Chromium(VI) Reduction by *Pseudomonas fluorescens* LB300 in Fixed-Film Bioreactor. *J Environ Eng* [Internet]. 1997 Aug [cited 2020 Jun 4];123(8):760–6. Available from: <http://ascelibrary.org/doi/10.1061/%28ASCE%290733-9372%281997%29123%3A8%28760%29>
  199. Naeem A, Batool R, Jamil N. Cr(VI) reduction by *Cellulosimicrobium* sp. isolated from tannery effluent. *Turk J Biol*. 2013;8.
  200. Quintelas C, Rocha Z, Silva B, Fonseca B, Figueiredo H, Tavares T. Removal of Cd(II), Cr(VI), Fe(III) and Ni(II) from aqueous solutions by an *E. coli* biofilm supported on kaolin. *Chem Eng J* [Internet]. 2009 Jul 1 [cited 2021 Sep 22];149(1–3):319–24. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1385894708007353>
  201. Husien S, Labena A, El-Belely E, Mahmoud H, Hamouda A. Application of *Nostoc* sp. for hexavalent chromium [Cr(VI)] removal: planktonic and biofilm. *Int J Environ Anal Chem* [Internet]. 2020 Jun 5 [cited 2021 Sep 22];1–22. Available from: <https://www.tandfonline.com/doi/full/10.1080/03067319.2020.1773454>
  202. Zinicovscaia I, Safonov A, Boldyrev K, Gundorina S, Yushin N, Petuhov O, et al. Selective metal removal from chromium-containing synthetic effluents using *Shewanella xiamenensis* biofilm supported on zeolite. *Environ Sci Pollut Res* [Internet]. 2020 Apr [cited 2021 Sep 22];27(10):10495–505. Available from: <http://link.springer.com/10.1007/s11356-020-07690-y>
  203. Hussain S, Quinn L, Li J, Casey E, Murphy CD. Simultaneous removal of malachite green and hexavalent chromium by *Cunninghamella elegans* biofilm in a semi-continuous system. *Int Biodeterior Biodegrad* [Internet]. 2017 Nov [cited 2021 Sep 22];125:142–9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830517308843>
  204. Dey S, Paul AK. Magnesium-induced biofilm development in *Arthrobacter* sp. SUK 1201 and removal of hexavalent chromium. *Soil Sediment Contam Int J* [Internet]. 2018 Jul 4 [cited 2021 Sep 22];27(5):383–92. Available from: <https://www.tandfonline.com/doi/full/10.1080/15320383.2018.1484688>
  205. Kumar H, Sinha SK, Goud VV, Das S. Removal of Cr(VI) by magnetic iron oxide nanoparticles synthesized from extracellular polymeric substances of chromium resistant acid-tolerant bacterium *Lysinibacillus sphaericus* RTA-01. *J Environ Health Sci Eng* [Internet]. 2019 Dec [cited 2021 Sep 22];17(2):1001–16. Available from: <http://link.springer.com/10.1007/s40201-019-00415-5>
  206. Tandon S, Jha M, Dudhwadkar S. Study on *Ochrobactrum pseudintermedium* ADV31 for the removal of hexavalent chromium through different immobilization techniques. *SN Appl Sci* [Internet]. 2020 Feb [cited 2021 Sep 22];2(2):296. Available from: <http://link.springer.com/10.1007/s42452-020-2103-y>
  207. Pratush A, Kumar A, Hu Z. Adverse effect of heavy metals (As, Pb, Hg, and Cr) on health and their bioremediation strategies: a review. *Int Microbiol* [Internet]. 2018 Sep [cited 2021 Sep 11];21(3):97–106. Available from: <http://link.springer.com/10.1007/s10123-018-0012-3>
  208. Azubuike CC, Chikere CB, Okpokwasili GC. Bioremediation techniques–classification based on site of application: principles, advantages, limitations and prospects. *World J Microbiol Biotechnol* [Internet]. 2016 Nov [cited 2021 Jul 26];32(11):180. Available from: <http://link.springer.com/10.1007/s11274-016-2137-x>
  209. Ahmad WA, Venil CK, Nkhalambayausi Chirwa EM, Wang Y-T, Sani MohdH, Samad AFA, et al. Bacterial Reduction of Cr(VI): Operational Challenges and Feasibility. *Curr Pollut Res* [Internet]. 2021 Jun [cited 2021 Sep 28];7(2):115–27. Available from: <https://link.springer.com/10.1007/s40726-021-00174-8>
  210. Apte A, Tare V, Bose P. Extent of oxidation of Cr(III) to Cr(VI) under various conditions pertaining to natural environment. *J Hazard Mater* [Internet]. 2006 Feb 6 [cited 2021 Sep 17];128(2–3):164–74. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389405004553>
  211. Lindsay DR, Farley KJ, Carbonaro RF. Oxidation of Cr(III) to Cr(VI) during chlorination of drinking water. *J Environ Monit* [Internet]. 2012 [cited 2021 Sep 17];14(7):1789. Available from: <http://xlink.rsc.org/?DOI=c2em00012a>
  212. Espinoza-Sánchez MA, Arévalo-Niño K, Quintero-Zapata I, Castro-González I, Almaguer-Cantú V. Cr(VI) adsorption from aqueous solution by fungal bioremediation based using *Rhizopus* sp. *J Environ Manage* [Internet]. 2019 Dec [cited 2021 Sep 28];251:109595. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0301479719313131>
  213. Benila Smily JRM, Sumithra PA. Optimization of Chromium Biosorption by Fungal Adsorbent, *Trichoderma* sp. BSCR02 and its Desorption Studies. *HAYATI J Biosci* [Internet]. 2017 Apr [cited 2021 Sep 28];24(2):65–71. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S197830191630434X>
  214. Xu X, Zhang Z, Huang Q, Chen W. Biosorption Performance of Multimetal Resistant Fungus *Penicillium chrysogenum* XJ-1 for Removal of Cu<sup>2+</sup> and Cr<sup>6+</sup> from Aqueous Solutions. *Geomicrobiol J* [Internet]. 2018 Jan 2 [cited 2021 Sep 28];35(1):40–9. Available from: <https://www.tandfonline.com/doi/full/10.1080/01490451.2017.1310331>
  215. Mondal NK, Samanta A, Roy P, Das B. Optimization study of adsorption parameters for removal of Cr(VI) using *Magnolia* leaf biomass by response surface methodology. *Sustain Water Resour Manag* [Internet]. 2019 Dec [cited 2021 Sep 28];5(4):1627–39. Available from: <http://link.springer.com/10.1007/s40899-019-00322-5>
  216. Vendruscolo F, da Rocha Ferreira GL, Antoniosi Filho NR. Biosorption of hexavalent chromium by microorganisms. *Int Biodeterior Biodegrad* [Internet]. 2017 Apr [cited 2020 Apr 9];119:87–95. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0964830516305054>
  217. Antony GS, Manna A, Baskaran S, Puhazhendi P, Ramchary A, Niraikulam A, et al. Non-enzymatic reduction of Cr (VI) and it's effective biosorption using heat-inactivated biomass: A



- fermentation waste material. *J Hazard Mater* [Internet]. 2020 Jun [cited 2021 Sep 28];392:122257. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0304389420302454>
218. Daneshvar E, Zarrinmehr MJ, Kousha M, Hashtjin AM, Saratale GD, Maiti A, et al. Hexavalent chromium removal from water by microalgal-based materials: Adsorption, desorption and recovery studies. *Bioresour Technol* [Internet]. 2019 Dec [cited 2021 Sep 28];293:122064. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960852419312945>
219. Sukumar C, Janaki V, Kamala-Kannan S, Shanthi K. Biosorption of chromium(VI) using *Bacillus subtilis* SS-1 isolated from soil samples of electroplating industry. *Clean Technol Environ Policy* [Internet]. 2014 Feb [cited 2021 Jul 16];16(2):405-13. Available from: <http://link.springer.com/10.1007/s10098-013-0636-0>
220. Sathvika T, Manasi, Rajesh V, Rajesh N. Adsorption of chromium supported with various column modelling studies through the synergistic influence of *Aspergillus* and cellulose. *J Environ Chem Eng* [Internet]. 2016 Sep [cited 2021 Sep 28];4(3):3193-204. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S221334371630238X>
221. Aravindhana R, Aafreen Fathima, Selvamurugan M, Raghava Rao J, Balachandran UN. Adsorption, desorption, and kinetic study on Cr(III) removal from aqueous solution using *Bacillus subtilis* biomass. *Clean Technol Environ Policy* [Internet]. 2012 Aug [cited 2021 Jul 19];14(4):727-35. Available from: <http://link.springer.com/10.1007/s10098-011-0440-7>